



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) Publication number:

0 337 589 B1

(12)

EUROPEAN PATENT SPECIFICATION

- (49) Date of publication of patent specification: **23.06.93** (51) Int. Cl.⁵: **C12N 15/30, C12P 21/02, A61K 39/012, C07K 13/00**
- (21) Application number: **89300328.5**
- (22) Date of filing: **13.01.89**

USSN: 10/ 723,123
Express Mail: EV 630723220 US
Attorney Docket: I-1998.407 US D2

- (54) **Recombinant and native group B eimeria tenella immunogens useful as coccidiosis vaccines.**

- (30) Priority: **15.01.88 US 145802**
22.12.88 US 286936
- (43) Date of publication of application:
18.10.89 Bulletin 89/42
- (45) Publication of the grant of the patent:
23.06.93 Bulletin 93/25
- (94) Designated Contracting States:
AT BE CH DE ES FR GB IT LI LU NL
- (56) References cited:
EP-A- 0 135 073
EP-A- 0 164 176
EP-A- 0 231 537
WO-A-86/00528
WO-A-88/06629

CHEMICAL ABSTRACTS, vol. 106, no. 17, 27th April 1987, pages 189-190, abstract no. 132966m, Columbus, Ohio, US; **L.E. CLARKE et al.**: "Isolation of lambdaamp3 genomic recombinants coding for antigens of Eimeria tenella"

- (73) Proprietor: **MERCK & CO. INC.**
126, East Lincoln Avenue P.O. Box 2000
Rahway New Jersey 07065-0900(US)
- (72) Inventor: **Liberator, Paul A.**
11 Bridge Court
Jackson New Jersey 08527(US)
Inventor: **Nollstadt, Karl H.**
91 Stonehenge Terrace
Clark New Jersey 07066(US)
Inventor: **Turner, Mervyn J.**
918 Boulevard
Westfield New Jersey 07090(US)
Inventor: **St. John Crane, Mark**
137 St. Paul Street
Westfield New Jersey 07090(US)
Inventor: **Karkhanis, Yashwant D.**
160 Coriell Avenue
Fanwood New Jersey 07023(US)
Inventor: **Chakraborty, Prasanta R.**
2242 Newark Avenue
Scotch Plains New Jersey 07076(US)
Inventor: **Profous-Juchelka**
39 Poplar Avenue,
Staten Island New York 10309(US)

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid (Art. 99(1) European patent convention).

CHEMICAL ABSTRACTS, vol. 109, no. 21, 21st November 1988, page 374, abstract no. 186706g, Columbus, Ohio, US; H. PROFOUS-JUCHELKA et al.: "Identification and characterization of cDNA clones encoding antigens of *Eimeria tenella*"

74 Representative: Cole, William Gwyn
European Patent Department, Merck & Co.,
Inc., Terlings Park, Eastwick Road
Harlow, Essex CM20 2QR (GB)

Description

BRIEF DESCRIPTION OF THE DRAWINGS

FIGURE 1 is a restriction map of group A clones.
 FIGURE 2 is a restriction map of group B clones.
 FIGURE 3 is a restriction map of group C clones.
 FIGURE 4 is a restriction map of group H clones.
 FIGURE 5 is a restriction map of group F clones.
 FIGURE 6 is a diagram of the pSC1N plasmid.
 FIGURE 7 illustrates the conversion of the CheY-ANF plasmid to the pJC264 plasmid.
 FIGURE 8 is a restriction map of the pJC264 plasmid.

BACKGROUND OF THE INVENTION

Coccidiosis is a disease caused by infection with one or more of the many species of coccidia, a subdivision of the phylum Protozoa. The coccidia are intracellular parasites which can infect a wide range of hosts and may result in severe economic loss to the sheep, goat, cattle, swine and poultry industry. Indeed, coccidiosis resulting from infection with *Eimeria* species has caused economically devastating losses to the poultry industry. Poultry is defined herein as domesticated birds that serve as a source of eggs or meat and that include among commercially important kinds chickens, turkeys, ducks, geese, guinea fowl, pheasants, pigeons and peafowl. Among domesticated birds, chicken production is the most susceptible to the economic losses from coccidiosis, although losses can also occur with turkeys, geese, ducks, and guinea fowl. Coccidiosis also produces serious losses in pheasants and quail raised in captivity. Coccidiosis may be acute and characterized by devastating flock mortality or the disease may be chronic and characterized by lack of weight gain.

Poultry are infected by coccidia following ingestion of the vegetative stage of the parasite, the sporulated oocyst. The infective stage, the sporozoite, is released in the intestine where it rapidly invades epithelial cells subsequently undergoing several generations of rapid intracellular asexual multiplication (schizogony) before entering the stage of sexual differentiation and mating (gametogony) leading to the formation of immature oocysts which are shed in the droppings and then undergo an extracellular sporulation process (sporogony) resulting in the generation of mature oocysts. Low level infection with any of the *Eimeria* species (spp.), *E. acervulina*, *E. mivati*, *E. mitis*, *E. praecox*, *E. hagani*, *E. necatrix*, *E. maxima*, *E. brunetti* and *E. tenella* results in a protective immunity to reinfection. There may be as many as twelve distinct cell types involved in the development of the parasite, each morphologically and antigenically different. At least three of these cell types have been shown to induce a protective immune response in the host, Rose and Hesketh, *Parasitol.* 73:25-37 (1976), McDonald et al., *Parasitol.* 93:1-7 (1986), Bhanushali and Long, In, *Research in Avian Coccidiosis*, Proc. of the Georgia Coccidiosis Conf., Athens, GA, USA pp. 526-534 (1986). Both the sporozoite as well as the first and second generation schizont appear to contain antigens which elicit an immunizing effect in chickens.

Unlike the sporozoite surface of other parasites such as *Plasmodium falciparum* which is composed of a single dominant antigen, Santoro et al., *J. Biol. Chem.* 258:3341-3345 (1983), the *Eimeria* spp., in particular, *E. tenella* sporozoite surface appears to be antigenically complex, Wisher, *Mol. Biochem. Parasitol.* 21:7-15 (1986). Because the sporozoite stage cannot be cultivated *in vitro* and large amounts of sporozoite material would be necessary for conventional biochemical analysis and for subunit vaccine evaluation, the purification of these antigens has posed a problem. A subunit vaccine as used herein is defined as a peptide, polypeptide or protein which is either isolated from one or more of the life stages of any species of *Eimeria* or is produced by recombinant DNA technology and which either individually or combined with other such peptides, polypeptides or proteins induces a protective immunity in poultry following vaccination. The recombinant antigens or immunogens will be the same as or similar to the peptides, polypeptides or proteins isolated from one or more life stages of *Eimeria*. Immunogen is defined as a substance that when introduced into the body stimulates an immune response which is protective in nature, such as the use of a vaccine to produce immunity against a microorganism. Immunity is defined as the non-susceptibility to the invasive or pathogenic effects of foreign organisms or the toxic effects of products of foreign organisms. The protective immunity may be either humoral or cell-mediated immunity. Humoral immunity is defined as specific immunity mediated by antibodies which are present in the plasma, lymph and tissue fluids of the body, and which may become attached to cells. Cell-mediated immunity is defined as specific immunity mediated by T lymphocytes. Antigen is used herein to define a substance capable of specifically combining

with specific antibody.

Solubilized E. tenella sporozoite proteins, identified by monoclonal antibodies prepared against intact E. tenella sporozoites, have been shown to protect chickens against challenge with infective oocysts. Schenkel et al., European Patent Application Number 135,712. Similar results were obtained with E. tenella merozoites prepared by the same techniques, Schenkel et al. European Patent Application Number 135,073. Immunogenic polypeptides have been isolated from E. tenella sporozoites, Murray and Galuska, U.S. Patent Number 4,639,372. There was no indication, however, that any individual polypeptide would protect chickens against E. tenella challenge.

Recombinant DNA technology has allowed for the identification of immunogenic Eimeria polypeptides and for the production of the polypeptides in sufficient quantities for vaccine development. Newman et al., European Patent Application 164,176, describe the isolation of a 25,000 dalton polypeptide from E. tenella which is made up of two subunits of 17,000 and 8,000 daltons respectively. The 25,000 dalton polypeptide has been produced by recombinant DNA technology utilizing a genomic DNA clone and has been shown to protect chickens against coccidiosis caused by E. tenella. Another immunogenic E. tenella polypeptide has been disclosed by Anderson and McCandliss, Patent Cooperation Treaty Application WO 86/00528. This peptide has been sequenced and is composed of 280 amino acids, has been produced by recombinant DNA technology utilizing both an oocyst genomic DNA clone and a clone isolated from total oocyst mRNA, and protects chickens against coccidiosis. Clark et al., Mol. and Biochem. Parasit. 22:79-87 (1987), recently disclosed the construction of genomic DNA expression libraries from E. tenella in Escherichia coli using the expression vector amp3. Clones expressing E. tenella immunogens were detected but none of the peptides were tested for immunogenic activity. Eimeria tenella sporozoite surface membranes have been labeled by various techniques to characterize potential surface immunogens, Wisher, Mol. Biochem. Parasit, 21:7-15 (1986). The major surface polypeptides which reacted with anti-E. tenella antibody were in the following ranges: 113-96 kD, 73-67 kD, 54-42 kD, 37-32 kD, and 18-14 kD.

SUMMARY OF THE INVENTION

Genes coding for novel Group B Eimeria tenella protein immunogens have been isolated and inserted into a novel expression vector which in turn has been used to transform appropriate hosts. The transformed host cells produce recombinant Group B E. tenella proteins which are capable of inducing immunity in chickens to coccidiosis. Antibody prepared against the recombinant protein immunogens is used to isolate and identify the native protein from disrupted E. tenella sporulated oocysts.

OBJECT OF THE INVENTION

It is accordingly, an object of the present invention to provide novel proteins of Eimeria tenella which can be used to immunize chickens against coccidiosis. Another object is to provide immunogenic proteins specifically associated with sporulated oocysts and sporozoites. A further object is to provide the deduced amino acid sequence of the immunogenic proteins. Another object is to isolate genes coding for the specific protein immunogens and to incorporate the genes into appropriate expression vectors. A further object is to transform an appropriate host with each of the recombinant vectors, to induce expression of the specific coccidial genes and to isolate the pure immunogens. Another object is to produce a novel expression vector for the expression of the specific coccidial proteins. A further object is to produce monospecific antibodies reactive against the immunogenic proteins.

DETAILED DESCRIPTION OF THE INVENTION

The present invention relates to coccidiosis vaccines based on either native or recombinant-derived purified protein immunogens and any microheterogeneous or subunit immunogen forms of the protein associated with sporulated oocysts, sporozoites, schizonts and merozoites of Eimeria tenella. Native protein as used herein refers to the full length protein produced by the appropriate Eimeria gene in the parasite. Recombinant-derived refers to the isolation of a gene for a desired protein and the use of that purified gene to construct a bacterium which will overproduce the desired protein. Subunit immunogen forms is defined as a portion of an immunogenic protein or polypeptide which has fewer amino acids than the native immunogenic moiety but contains the immunogenic site or sites of the immunogen. Microheterogeneous forms as used herein refers to a single gene product, that is a protein produced from a single gene unit of DNA, which is structurally modified following translation. These structural modifications, however, do not result in any significant alterations of the immunogenic activity of the protein. The modifications may take

place either in vivo, in the parasite, or during the isolation and purification process. In vivo modification may result in, but is not limited to, acetylation at the N-terminus, proteolysis, glycosylation or phosphorylation. Proteolysis may include exoproteolysis wherein one or more terminal amino acids are sequentially, enzymatically cleaved to produce microheterogeneous forms which have fewer amino acids than the original gene product. Proteolysis may also include endoproteolytic modification that results from the action of endoproteases which cleave the peptide at specific locations within the amino acid sequence. Similar modifications can occur during the purification process which may result in the production of microheterogeneous forms. The most common modification occurring during purification is proteolysis which is generally held to a minimum by the use of protease inhibitors.

The invention further relates to isolation and purification of the genetic information responsible for individual protein and the methods of expressing the corresponding immunogenic proteins. Polypeptide or protein as used herein refers to a linear polymer of amino acids bound together with amide linkages. The sequence of amino acids in the chain is of critical importance in the biological functioning of the protein or polypeptide. Polypeptide and protein are used interchangeably herein. Immunogen as used herein refers to molecules or macromolecules which when introduced into an animal body stimulates a humoral and/or a cellular immune response which is functional in nature, that is an immunity which protects the animal from a specific infection. In the instant case an immunogen will produce an immune response, either humoral, cellular or both which will protect poultry against infection with *Eimeria* species which cause coccidiosis.

Eimeria tenella oocysts are isolated from the cecal contents of chickens infected 4 to 10 days earlier, preferably 7 days, while *E. acervulina* oocysts are isolated from feces and intestinal contents of chickens infected 5 to 6 days earlier. The cecal contents and feces are individually physically disrupted in a Waring Blender, in distilled water and digested with a proteolytic enzyme, preferably pepsin. Debris and pepsin are removed by centrifugation in distilled water. A partially pure oocyst fraction is collected by flotation in about 2.2 M sucrose, Jackson, Parasitol. 54:87-93 (1964), and further treated by incubation in sodium hypochlorite at a concentration of about 5 to about 6 percent, preferably 5.25%, in water at about 4°C for approximately 10 minutes. The sodium hypochlorite is removed by several washes in sterile phosphate buffered saline (PBS) at about pH 7.6 to obtain purified, sterile oocysts. Oocysts are allowed to sporulate in a shaking water bath for about 48 hours at about 20°C, Edgar, Trans. Am. Micr. Soc. 62:237-242 (1954).

Sporulated oocysts are suspended in PBS and disrupted in a Bransonic cell disruptor (Branson), with a tapered probe at about 0°C. Sonication is carried out with short bursts, about 30 seconds, to prevent overheating, with 90 percent breakage occurring within about 5 to about 20 minutes. A detergent is added to the sonicate, preferably Zwittergent 3-12 (Calbiochem) about 0.1% w/v and the mixture is stirred at about 4°C for about 18 hours. The detergent treated sporulated oocyst preparation is centrifuged at about 27,000 x g for about 30 minutes and the supernatant fluid collected.

Sporozoites are prepared by grinding a suspension of purified sporulated oocysts, about 5×10^7 /ml in PBS, at about pH 7.6, at about 500 rpm for about 5 minutes at about 4°C in a tissue homogenizer with a loose-fitting pestle following the procedure of Patton, Science 150:767-769 (1965). The *E. tenella* disrupted material is collected by centrifugation. The pellet consists of unbroken oocysts, sporocysts and oocyst shells which is resuspended in an excysting solution containing about 0.25% (w/v) trypsin and about 4% (w/v) taurodeoxycholic acid (Sigma) in a buffered solution such as Hanks balanced salt solution (pH 7.4). The *E. acervulina* pellet, also composed of unbroken oocysts, sporocysts and oocyst shells, was resuspended in an excysting solution containing about 0.125% (w/v) trypsin (1:250) and about 1.0% taurodeoxycholic acid in a buffered solution such as Hank's Balanced salt solution (pH 7.4). The resuspended pellets are incubated at about 41°C in an atmosphere containing about 5% CO₂. Excysting was allowed to continue for about 1/2 hour for *E. acervulina* and about 1 hour for *E. tenella* after which time the solutions are removed by centrifugation. Sporozoites are isolated using a DE-52 anion exchange column employing the method of Schmatz et al. J. Protozool. 31:181-183 (1984). Purified sporozoites are disrupted by freezing and thawing at least 3 times, and sonicated until disrupted in PBS containing about 1mM phenylmethylsulfonylfluoride.

Both the sporulated oocyst and the sporozoite cell free preparations are separated by gel permeation chromatography, preferably Sephadex S-200 (Pharmacia) in a separation buffer containing about 50 mM Na₂HPO₄-NaH₂PO₄, pH about 7.2 and about 0.1% Zwittergent 3-12. Each preparation is added to the column, about 8 x 44 cm and eluted with the separation buffer. Elution is monitored by absorbance at 230 nm and the fractions, about 14 ml per fraction, collected. The fractions are analyzed by linear gradient sodium dodecyl sulfate (SDS) polyacrylamide gel electrophoresis (SDS-PAGE) and the fractions pooled according to these profiles. Pooled fractions were dialyzed against a bicarbonate buffer and tested for their ability to protect chickens against challenge with infective *E. tenella* sporulated oocysts. Two day old broiler pullets are immunized intramuscularly with pooled fractions of sporulated oocyst or sporozoite cell free immunogens, about 5 µg to about 50 µg protein in PBS. The cell free immunogen is precipitated to alum

(about 0.4% final concentration) in a total volume of about 0.12 ml per dose per bird. The alum-immunogen precipitation complex is prepared by the technique of Weir, Handbook of Experimental Immunology, Blackwell Scientific Publications, London pg. A3.11 (1978). Immunization was repeated at days nine and sixteen and the birds are challenged on day 23, seven days after the final immunization, with infective *E. tenella* sporulated oocysts. A single fraction from each preparation protected the chickens from sporozoite challenge. These fractions had similar elution and electrophoresis profiles suggesting that the polypeptides may be similar. The most active immunogenic fraction isolated from sporulated oocysts is found in column fractions 84-94 and is designated Fraction V.

Antiserum is produced against the immunoprotective fractions of *Eimeria tenella* sporulated oocysts (Fraction V), sporozoites, sonicated unsporulated oocysts, second generation schizonts and *E. acervulina* sonicated sporozoites. The *E. tenella* schizonts are prepared from chicken intestinal cells about four days post-infection according to the protocol of James, Parasitol. 80:301-312 (1980). Blood is collected from the antibody producing animals, preferably rabbits, prior to initiation of the immunization procedure and the preimmune serum is isolated and stored for control purposes. The rabbits are given multiple immunization injections with one of the above described immunogens, about 20 µg to about 80 µg of protein per immunization. The initial immunization is given with an acceptable adjuvant, generally equal volumes of immunogen and adjuvant. Acceptable adjuvants include Freund's complete, Freund's incomplete, alum-precipitate, water-in-oil emulsion containing *Corynebacterium parvum* and tRNA, with Freund's complete adjuvant being preferred for the initial immunization. Freund's incomplete adjuvant is preferred for all booster immunizations. The initial immunization consists of the administration of about 1 ml of emulsion at multiple subcutaneous sites on the backs of the rabbits. Booster immunizations utilizing an equal volume of immunogen are given at about one month intervals and are continued until adequate levels of antibodies are present in an individual rabbits serum. Blood is collected and serum isolated by methods known in the art. The anti-coccidial antisera are characterized by serological analysis, preferably Western blot analysis using antigens obtained from unsporulated oocysts, sporulated oocysts, sporozoites and schizonts. Antigen as used herein is defined as any substance that can combine with an antibody. Immunogens as described above are considered antigens when used to characterize the specific antibody.

The parasite immunogens to be used for Western blot analysis, about 50 µg, as described above, are mixed in about equal volumes with about 2x concentrated sample buffer consisting of about 0.1 M Tris HCl, about pH 6.8, about 4% sodium dodecyl sulfate (SDS), about 20% (v/v) glycerol, about 10% (v/v) 2-mercaptoethanol, and about 0.002% (v/v) bromophenol blue. The samples are boiled for about 3 minutes and electrophoresed on a 5-20% linear gradient of polyacrylamide gel (PAGE) containing SDS by the method of Laemmli, Nature 227:680-684 (1970). The proteins separated by SDS-PAGE are electrophoretically transferred to nitrocellulose by the method of Towbin et al., Proc. Natl. Acad. Sci. USA 76:4350-4354 (1979), and the nitrocellulose is blocked with 0.5% gelatin in phosphate buffered saline about pH 7.4. The blocked nitrocellulose is incubated overnight at room temperature in about 20 ml of the appropriate antiserum diluted about 1:5 to 1:400 in TEN buffer (about 50 mM Tris-HCl, about 150 mM NaCl and about 5 mM ethylenediamine tetraacetic acid (EDTA) at a pH of about 7.4) containing about 0.25% gelatin and 0.05% Triton X-100. Bound antibody is detected by the addition of ¹²⁵I-protein A.

Since none of the coccidial polypeptides, described above, which confer immunity, are capable of being purified to homogeneity by known separation or purification methods it has been impossible to characterize the amino acid composition of the individual polypeptides. Consequently, the antibodies directed against the various *Eimeria* antigens are used to identify, by immunological methods, protective coccidial immunogenic polypeptides produced by recombinant DNA technology. Recombinant DNA technology is defined herein as technology which allows segments of genetic information, DNA, from different cells, usually from different organisms, to be joined end-to-end outside the organisms from which the DNA was obtained and to incorporate this hybrid DNA into a cell that will allow the production of the protein for which the original DNA encodes. Genetic information, DNA or mRNA, is isolated from sporulating oocysts or sporozoites, incorporated into an appropriate cloning vector, transduced into an appropriate host cell and products of the host cell screened for the production of polypeptides which bind to the anti-*E. tenella* antibodies. The identified genes expressing the immunoreactive polypeptides are incorporated into an appropriate expression vector and expressed in an appropriate host cell system.

Cloning vector as used herein is defined as a DNA sequence which allows the incorporation of specific experimental foreign DNA, with the combined DNA being introduced into a host cell that can exist in a stable manner and express the protein dictated by the experimental DNA. The foreign DNA combined with the vector DNA constitutes a recombinant DNA molecule which is derived from recombinant technology. Cloning vectors may include plasmids, bacteriophage, viruses and cosmids. It is to be understood that any cloning vector may be used to clone the novel *Eimeria* immunogen DNA sequences, with the lambda gt11

bbeing preferred. Host cells for cloning, DNA processing and initial expression generally include bacteria. The preferred cloning host is *Escherichia coli*. Expression vectors are defined herein as DNA sequences that are required for the transcription of cloned copies of genes and the translation of their mRNAs in an appropriate host. Such vectors can be used to express either procaryotic or eucaryotic genes in a variety of hosts such as bacteria, blue-green algae, yeast cells, insect cells and animal cells. The immunogens may also be expressed in a number of virus systems. Specifically designed vectors allow the shuttling of DNA between bacteria-yeast or bacteria-animal cells. An appropriately constructed expression vector should contain: an origin of replication for autonomous replication in host cells, selective markers, a limited number of useful restriction enzyme sites, a high copy number, and strong promoters. A promoter is defined as a DNA sequence that directs RNA polymerase to bind to DNA and to initiate RNA synthesis. A strong promoter is one which causes mRNAs to be initiated at high frequency. Expression vectors may include, but are not limited to, cloning vectors, modified cloning vectors, specifically designed plasmids or viruses.

The unique immunogenic proteins of the present invention may exist as, but are not limited to, the complete proteins specified by the defined gene in *Eimeria*, native protein or as any fragment or subunit thereof, or as hybrids of the complete protein or its fragments or subunits. The complete protein, as used herein, refers to the full length polypeptide produced by the appropriate *Eimeria* gene. The complete protein may be obtained by purification from the appropriate species of *Eimeria*, or by expression in an appropriate expression vector of the corresponding recombinant derived gene product. Protein fragments or subunits refers to any portion of the protein which contains fewer amino acids than the complete protein and retains the ability to induce anticoccidial immunity. Hybrid proteins include, but are not limited to, fusion proteins or proteins resulting from the expression of multiple genes within the expression vector. A fusion protein is defined as one in which a limited number of amino acids coded for by the expression vector are expressed and the expression results in their attachment to the specific immunogenic polypeptide. Proteins resulting from multiple genes may include the specific immunogenic polypeptide linked to a second polypeptide or peptides by peptide bonds that enhance immune reactivity. The enhancing polypeptide portion may have the capability of increasing the immune response to the coccidial immunogen.

The appropriate coccidial DNA is isolated and identified by reacting the gene derived protein with anti-Fraction V and anti-sporozoite antibodies. Recombinant coccidial polypeptides are produced by cloning the natural gene from either genomic DNA or cDNA. Genomic DNA, a preferred method of obtaining specific genes, is extracted from sporocysts or sporozoites by disrupting the parasites, about 1.5×10^8 , by treatment with about 0.5% SDS and about 15 mM EDTA. The released DNA is solubilized by digestion with a proteolytic enzyme, preferably Proteinase K, about 100 $\mu\text{g/ml}$ at about 50°C for about 3 hours. Genomic DNA is purified by about two extractions with phenol, about two extractions with a mixture of phenol, chloroform and isoamyl alcohol (about 25:24:1), about two extractions with chloroform and isoamyl alcohol (about 24:1) and about two successive precipitations with sodium acetate/ethanol. The DNA is washed twice with about 70% ethanol and resuspended in Tris-HCl, about 10 mM and EDTA, about 1 mM (TE) at the approximate concentration of about 5×10^8 parasite equivalents per ml. Any associated RNA is selectively removed by digestion with RNase, preferably heat inactivated RNase A, at a concentration of about 50 $\mu\text{g/ml}$ for about 60 minutes at about 37°C . The RNase A and any other residual proteins are removed by a secondary digestion with Proteinase K in about 0.5% SDS/15 mM EDTA for about 3 hours at about 50°C . The genomic DNA is then extracted with organic solvents, precipitated with ethanol and washed with about 70% ethanol and collected by centrifugation. The genomic DNA pellet is suspended in TE at a concentration of about $2-3 \times 10^9$ sporozoite equivalents/ml and quantitated by absorbance at 260 nm. Coccidial DNA is prepared for cloning by either physical, Old and Primrose, Principles of Gene Manipulation, 2nd Ed. University of California Press, p. 20 (1981) or chemical, Smithies et al., Science 202:1284-1289 (1978), fragmentation of high molecular weight DNA. The genomic DNA is then incorporated into an appropriate cloning vector, see cloning vectors for cDNA below. The cloning vectors are transduced into a host cell and screened by a procedure similar to that of Huynh et al., In "DNA cloning: A practical approach", Vol. I, Glover Ed., IRL Press Oxford, pp. 49-78 (1985). Positive clones are transferred to expression vectors engineered for high volume production of the desired immunogenic protein. The expression vectors, see below, are transformed into suitable host cells, see below, for the production of immunogenic protein.

The most preferred process for obtaining genetic information for the production of coccidial immunogenic polypeptides is the isolation of mRNA coding for a specific protein. Total RNA is isolated from oocysts, sporulated for about seven hours, and sporozoites using the guanidinium thiocyanate method of Chirgwin et al., Biochem. 18:5294-5299 (1979). Polyadenylated RNA is selected by oligo (dT)-cellulose chromatography, Aviv and Leder, Proc. Nat. Acad. Sci. USA 69:1408-1412 (1972). Utilizing the polyadenylated RNA, about 6 to about 9 μg , first and second-strand cDNA reactions are performed using a reverse transcriptase such as AMV-reverse transcriptase, a RNase such as RNase H and a DNA polymerase such as DNA polymerase I

following the procedure described by Gubler and Hoffman, Gene 25:263-269 (1983). The cDNA is methylated with a methylase such as Eco RI methylase, blunt-ended with a polymerase such as T4 DNA polymerase and ligated to phosphorylated oligonucleotide linkers such as Eco RI dextranucleotide linkers with a DNA ligase such as T4 DNA ligase. The linker ligated cDNAs are digested to completion with a restriction enzyme such as EcoRI and the digested linkers removed by repeated precipitations with absolute ethanol out of 2 M NH₄-acetate, Okayama and Berg, Mol. Cell. Biol. 2:161-170 (1982). The cDNA was further purified on an Elutip-d column (Schleicher & Schell). Restriction enzymes or restriction endonucleases are enzymes that recognize specific nucleotide base sequences within double-stranded DNA and cleave the two strands at a specific location within the recognition sequence. The purified cDNA, about 100 ng to about 500 ng, with 300 ng being preferred, is ligated into about 7.5 µg of commercially purchased, EcoRI-digested, alkaline phosphatase treated γgt11 vector DNA and packaged in vitro with commercially available packaging extracts according to the manufacturer's instructions (Amersham). Other acceptable vectors can be used, but γgt11 is preferred because it allows the inducible expression of Eimeria antigens in E. coli as β-galactosidase fusion proteins. Aliquots of the packaged phase are transduced into Escherichia coli host strain Y1088 and these are plated on Luria-Bertani (LB) medium agar plates using about 2.5 ml LB soft agar containing about 600 µg/ml X-gal (5-bromo-4-chloro-3-indolyl-β-D-galactopyranoside) and about 16 mM isopropyl-β-D-thiogalactopyranoside (IPTG).

A cDNA library consisting of approximately 1×10^7 independent recombinant phage clones is generated. The nonrecombinant background, as determined by growth on X-gal/IPTG plates, is estimated to be about 13%.

Screening of the cDNA library is accomplished by the method of Huynh, et al. "In: DNA Cloning: A Practical Approach", Vol. I, Glover, Ed., IRL Press, Oxford, pp. 49-78 (1985). Packaged phage from the unamplified cDNA library were transduced into E. coli strain Y1090 as described by Huynh, supra, and plated at an appropriate density, about 0.5 to about 1.0×10^5 plaque forming units (pfu) per plate. The plates are incubated, at about 42°C for about 3 hours, overlaid with nitrocellulose filters presoaked in about 10 mM IPTG, and reincubated overnight at about 37°C. The filters are removed, blocked with about 20% fetal calf serum in an acceptable buffer, such as Tris buffered saline (TBS) (about 50 mM Tris-HCl about 150 mM NaCl, at a pH of about 8.0) containing about 0.05% Tween 20 (TBST), and incubated with the appropriate antibody, generally rabbit anti-sporozoite antibody or rabbit anti-Fraction V antibody, diluted about 1:100 in TBST containing about 20% fetal calf serum for an appropriate length of time. All antisera are exhaustively preabsorbed with a concentrated lysate of lambda gtl1 lysogen BNN93. Antibody binding sites are detected by contacting the filters with ¹²⁵I-protein A. Positive plaques are picked, replated, and rescreened until each clone is shown to be plaque pure. An initial screen of the sporulated oocyst library of about 1×10^6 independent recombinants with rabbit anti-sporozoite antibody results in the isolation of about 57 antigen expressing phage. Secondary and tertiary rescreening reveals that greater than 29% of the clones initially identified remain positive.

Cross-screening involves the spotting of about 1 µl of phage lysate from each plaque purified clone on a lawn of E. coli Y1090 cells with recombinant fusion proteins being induced as previously described. The proteins are transferred to nitrocellulose and immunoblotted as described above. The cross-screening antisera include rabbit anti-E. tenella unsporulated oocyst antibody, rabbit anti-E. tenella sporozoite antibody, rabbit anti-Fraction V and rabbit anti-E. tenella schizont antibody. All antisera are exhaustively preabsorbed with a concentrated lysate of γgt11 lysogen BNN93.

Recombinant and wild type γgt11 phage are introduced as lysogens into E. coli host strain Y1089 at a multiplicity of about 10. Lysogenized clones are grown in about 10 ml of Luria-Bertani (LB) medium supplemented with about 50 µg/ml ampicillin at about 32°C until an optical density at 600 nm of 0.25 is reached. Phage replication is induced by a temperature shift to about 45°C for about 20 minutes and the synthesis of β-galactosidase fusion proteins is induced by the addition of about 10 mM IPTG to the culture medium. The cells are incubated and collected by centrifugation and the pellets are resuspended in about 250 µl of NET buffer, about 50 mM Tris-HCl, pH about 7.5, about 150 mM NaCl, about 5 mM ethylenediaminetetraacetic acid (EDTA), with about 2% SDS. The cells are lysed by boiling and the bacterial DNA is removed by centrifugation. The supernatant fluids are analyzed on about 5% SDS-PAGE under denaturing conditions. Duplicate gels are run with one being stained with silver stain (Biorad) and the other immunoblotted by the method of Towbin et al., Proc. Natl. Acad. Sci. USA 76:4350-4354 (1979).

Monospecific antibodies to each of the recombinant immunogens are affinity purified from polyclonal antisera by a modification of the method of Hall et al., Nature 311:379-382 (1984), prepared by immunizing rabbits as described above with purified recombinant E. tenella proteins as described below or prepared as monoclonal antibodies using the technique of Kohler and Milstein, Nature 256:495-497 (1975). Monospecific antibody as used herein is defined as a single antibody species or multiple antibody species with

homogeneous binding characteristics for the relevant antigen. Homogenous binding as used herein refers to the ability of the antibody species to bind to a specific antigen or epitope, such as those associated with the specific native or recombinant *E. tenella* group immunogens. The Hall technique of preparing monospecific antibodies from polyclonal antiserum requires the preparation of filter plaque lifts from purified recombinant clones as is done for screening. Approximately 2×10^5 plaque forming units are plated to give close to semiconfluent lysis at the end of the 37°C incubation period. The nitrocellulose is removed from the plates and is blocked with about 20% fetal calf serum in TBST for about 4 hours and incubated overnight with about 20 ml of the preabsorbed polyspecific serum, diluted about 1:200 with about 20% fetal calf serum in TBST containing about 0.02% NaN_3 . The filters are washed at least 5 times with about 50 ml TBST for at least 20 minutes and 1 time with about 0.15 mM NaCl and about 0.05% Tween 20. The antibodies are eluted with an acceptable eluant, such as about 0.2 M glycine-HCl, about 0.15 M NaCl and about 0.05% Tween 20, at a pH of about 2.8 for about 30 minutes. The pH is adjusted to about 8.0 and the antibodies are stored.

Monoclonal antibody reactive against each of the recombinant *E. tenella* group immunogens, antigens or epitopes is prepared by immunizing inbred mice, preferably Balb/c with the appropriate recombinant protein. The mice are immunized intraperitoneally with about 100 ng to about 10 μg , preferably about 1 μg recombinant immunogen per 0.5 ml in an equal volume of an acceptable adjuvant. Such acceptable adjuvants include, but are not limited to, Freund's complete, Freund's incomplete, alum-precipitate, water-in-oil emulsion containing *Corynebacterium parvum* and tRNA. The mice are given intravenous booster immunizations of an equal amount of recombinant immunogen without adjuvant at about days 14, 21, and 63 post primary immunization. At about day three after the final booster immunization individual mice are serologically tested for anti-recombinant immunogen antibody. Spleen cells from antibody producing mice are isolated and fused with murine myeloma cells, such as SP-2/0 or the like, by techniques known to the art, see Kohler and Milstein, Nature 256: 495-497 (1975). Hybridoma cells are selected by growth in hypoxanthine, thymidine and aminopterin in an appropriate cell culture medium such as Dulbecco's modified Eagle's medium (DMEM). Antibody producing hybridomas are cloned, preferably using the soft agar technique of MacPherson, Soft Agar Techniques, in Tissue Culture Methods and Applications, Kruse and Paterson, Eds. Academic Press, p. 276 (1973). Discrete colonies are transferred into individual wells of culture plates for cultivation in an appropriate culture medium. Antibody producing cells are identified by screening with the appropriate *E. tenella* recombinant immunogen. Immunogen positive hybridoma cells are maintained by techniques known in the art. Specific anti-recombinant *E. tenella* monoclonal antibodies are produced by cultivating the hybridomas in vitro or preparing ascites fluid in mice following hybridoma injection by procedures known in the art.

The parasite antigens are assayed by Western Blot analysis as described above. The clones of interest may be placed into four antigenic groups, according to the reaction of the expressed polypeptides with the above described antisera, see Table 1. Different clones of the same group express portions of the same polypeptide, as judged by antibody reactivity, DNA crosshybridization, and restriction endonuclease mapping, see below.

TABLE 1

IMMUNE REACTIVITY OF ISOLATED CLONE PRODUCTS					
CLONE	ANTI FRACTION V	ANTI-E.t. UNSPORULATED OOCYST	ANTI-E.t. SPOROZOITE	ANTI-E.t. SCHIZONT	ANTI-E.a. SPOROZOITE
A	+	+	+	-	+
B	+	-	+	-	+
C	+	-	+	-	-
H	+	-	+	n.d.	-
F	+	n.d.	n.d.	n.d.	n.d.

E.t. denotes *Eimeria tenella* while E.a. denotes *Eimeria acervulina*. A (+) denotes that the antibody can react with the specific recombinant derived protein while a (-) denotes a lack of such a response and n.d. means not done.

Purification of the cDNA inserts from γ t11 clones is accomplished by cutting the recombinant phage DNA to completion with EcoRI, about five fold enzyme excess, in a reaction buffer composed of about 50

Sporozoites were freeze-thawed 3 times (dry ice to room temperature and sonicated until disrupted in PBS with 1 mM phenylmethylsulfonylfluoride as protease inhibitor to provide sporozoite antigen. Protein concentrations were determined by the method of Lowry *et al.*, J. Biol. Chem. 193: 265-275, 1951 and antigens were stored in liquid N₂.

EXAMPLE 2

Production of Anti-*Eimeria tenella* Unsporulated, Oocyst, Sporulated Oocyst, Sporozoite, Schizont, Anti-Fraction V and Anti-*Eimeria acervulina* Sporozoite Antibodies

Rabbits (New Zealand White, female) were multiply immunized with one of the various immunogens described in Example 1. Each immunization dose contained 50 µg of protein. The first immunization was given in Freund's complete adjuvant. Subsequent immunizations were given in Freund's incomplete adjuvant. The antigen adjuvant mixture was prepared by emulsifying 0.5 ml of antigen containing 50 µg protein in PBS with 0.5 ml of adjuvant. One ml of emulsion was then administered subcutaneously in multiple sites on a shaved area of the rabbit back. Secondary booster immunizations were given at approximately one month intervals following primary immunization. Animals were bled and immune sera prepared at approximately monthly intervals, starting six weeks after the start of the immunization schedule. Immune activity and specificity was determined by Western blot analysis using the specific extract antigens from Example 1 and the technique of Towbin *et al.*, Proc. Natl. Acad. Sci. USA 76:4350-4354 (1979). Each antibody was specific for its corresponding immunogen, antigen.

EXAMPLE 3

Immunization of Two-Day-Old Chickens Against Coccidiosis with Fraction V Immunogens

Broiler chicks were immunized with Fraction V immunogen as described in Example 1. The dosage was based on protein content as determined by the method of Lowry *et al.*, J. Biol. Chem. 193: 265-275 (1951) and was given intramuscularly on days 2, 9 and 16 following hatching. Experimental and control chickens were challenged one week after the last immunization with an oral inoculation of 5×10^3 *E. tenella* oocysts. Six days after challenge the chickens were killed and the severity of the lesions in the ceca were determined according to the method of Johnson and Reid, Exp. Parasitol, 28: 30-36 (1970).

The following results were obtained.

TABLE 4

Immunogen	Dose (µg)	Number of Birds	Mean Group Lesion Score
Fraction V	10.0	8	1.0
Fraction V	1.0	8	1.6
Fraction V	0.10	8	2.9
None	-	8	3.4

These results show that Fraction V immunogen can be used to immunize two-day-old chickens. An intramuscular inoculation provides a high level of protection against the disease as indicated by the absence of severe lesion development in immune birds after a normally virulent infection.

TABLE 2
Amino Acid and Nucleotide Sequences
of the CheY Protein

5		10	20	30	40	50
		*	*	*	*	*
10	ATG GCG GAT AAA GAA CTT AAA TTT TTG GTT GTG GAT GAC TTT TCC ACC ATG CGA					
	MET ALA ASP LYS GLU LEU LYS PHE LEU VAL VAL ASP ASP PHE SER THR MET ARG					
		10				
15		60	70	80	90	100
		*	*	*	*	*
20	CGC ATA GTG CGT AAC CTG CTG AAA GAG CTG GGA TTC AAT AAT GTT GAG GAA GCG					
	ARG ILE VAL ARG ASN LEU LEU LYS GLU LEU GLY PHE ASN ASN VAL GLU GLU ALA					
		20		30		
25	110	120	130	140	150	160
	*	*	*	*	*	*
30	GAA GAT GGC GTC GAC GCT CTC AAT AAG TTG CAG GCA GGC GGT TAT GGA TTT GTT					
	GLU ASP GLY VAL ASP ALA LEU ASN LYS LEU GLN ALA GLY GLY TYR GLY PHE VAL					
		40		50		
35	170	180	190	200	210	
	*	*	*	*	*	
40	ATC TCC GAC TGG AAC ATG CCC AAC ATG GAT GGC CTG GAA TTG CTG AAA ACA ATT					
	ILE SER ASP TRP ASN MET PRO ASN MET ASP GLY LEU GLU LEU LEU LYS THR ILE					
		60		70		
45	220	230	240	250	260	
	*	*	*	*	*	
50	CGT GCG GAT GGC GCG ATG TCG GCA TTG CCA GTG TTA ATG GTG ACT GCA					
	ARG ALA ASP GLY ALA MET SER ALA LEU PRO VAL LEU MET VAL THR ALA					
		80				

Linker amino acids are defined herein as those amino acids used to link an E. tenella defined gene, one which produces a native protein, to a fusion protein. Any amino acid or group of amino acids may be used as linkers, however, the preferred amino acid sequence and nucleotide sequence of the peptide linking the CheY protein to the E. tenella protein is:

acceptable adjuvant. Acceptable adjuvants include, but are not limited to, Freund's complete, Freund's incomplete, double emulsions, anhydrous oils, alum-precipitate, water-in-oil emulsion containing *Corynebacterium parvum* and t-RNA. The preferred adjuvant is alum-precipitate, in which the immunogen has been precipitated with aluminum hydroxide such as Alhydrogel™. Immunization of chickens with recombinant derived *E. tenella* immunogens results in immunity to coccidiosis. Protective immunity is achieved by administration of from about 1.0 ng to about 100 µg, with about 100 ng to about 10 µg being preferred.

The following examples illustrate the present invention without, however, limiting the same thereto.

EXAMPLE 1

Preparation of Oocysts, Sporulated Oocysts, Sporozoites and Schizonts and the Corresponding Immunogens and Antigens

Eimeria tenella oocysts were isolated from cecal cores (coalesced masses of oocysts) from chickens infected 7 days earlier. *Eimeria acervulina* oocysts were isolated from feces and intestinal contents of chickens infected 5 to 6 days earlier. The isolated cecal cores and feces were separately disrupted in a Waring Blender (in distilled water), digested with pepsin (2 mg/ml) at pH 2.0 at 39°C for 1 hour. Large amounts of debris and the pepsin were removed from pelleted material after centrifugation (1,000 x g) in distilled water. A partially pure oocyst fraction was isolated from the pellet by flotation in 2.2 M sucrose (Jackson, Parasitol, 54: 87-93, 1964) and this crude material was further treated by incubating in cold Clorox (5.25% sodium hypochlorite, at 4°C) for 10 minutes. The sodium hypochlorite was removed by several washes in sterile phosphate-buffered saline (PBS) pH 7.6 to obtain purified and sterile oocysts. Oocysts were sporulated in a shaking water bath at 20°C for 48 hours (Edgar, Trans. Am. Micr. Soc. 62: 237-242, 1954). Sporulated oocysts were stored in PBS (pH 7.6) at 4°C.

Fully sporulated oocysts were sonicated on ice in a Bransonic cell disruptor, with a tapered probe. Sonication was performed using a 30 second on/off cycle to prevent overheating. Following this procedure, 90% breakage was achieved within 10-15 minutes. Detergent (Zwittergent 3-12, Calbiochem, 0.1% w/v) was added, and the mixture was stirred at 4°C for 18 hours. After centrifugation at 27,000 x g for 30 minutes, the supernatant was subjected to gel permeation chromatography on Sephadex S-200 (Pharmacia).

A column of Sephadex S-200 (8 x 44 cm) was equilibrated at 4°C with 50 mM Na₂HPO₄-NaH₂PO₄, pH 7.2 and 0.1% Zwittergent 3-12. The sonicate was applied to the column, eluted with the same buffer and fractions collected (14 ml) and monitored by absorbance at 230 nm. Fractions were pooled according to the SDS-PAGE profile. Pooled fractions were dialysed against eight liters of 10 mM ammonium bicarbonate at 4°C for one week with three changes of buffer, and were then freeze-dried. The lyophilized fractions were dissolved in glass-distilled water and were tested for *in vivo* activity, chicken protection. *In vivo* activity was routinely found between fractions 84-94. The protective *Eimeria tenella* fractions were pooled and designated Fraction V. For some batches, S-200 chromatography was performed in 50 mM ammonium bicarbonate, pH 7.7, containing 0.05% Zwittergent. This had no effect on the elution profile or on the *in vivo* efficacy.

Second generation schizonts were prepared from chicken intestinal cells four days post-infection according to the protocol of James, Parasitol, 80: 301-312 (1980).

Immunogens for antibody production were prepared as follows. A 2 ml suspension of purified sporulated oocysts (5 x 10⁷ per ml PBS, pH 7.6) was ground at 500 rpm for 5 minutes at 4°C in a tissue homogenizer with a loose-fitting pestle (Patton, Science 150: 767-760, 1965) and the supernatant fluid resulting from the disruption of the oocysts was removed after centrifugation (600 x g for 10 minutes). The *E. tenella* pellet, composed of unbroken oocysts, sporocysts, and oocyst shells, was resuspended in an excysting solution containing 0.25% (w/v) trypsin (1:250) and 4.0% (w/v) taurodeoxycholic acid (Sigma) in Hanks balanced salt solution (pH 7.4) and incubated at 41°C in 5% CO₂ (Patton et al, J. Parasitol. 65: 526-530, 1979). The *E. acervulina* pellet, also composed of unbroken oocysts, sporocysts and oocyst shells, was resuspended in an excysting solution containing 0.125% (w/v) trypsin (1:250) and 1.0% taurodeoxycholic acid in Hank's Balanced salt solution (pH 7.4) the pellets were incubated at 41°C in an atmosphere containing 5% CO₂. Excystation was allowed to continue for 1/2 hour for *E. acervulina* and 1 hour for *E. tenella* after which the excysting solution was removed by centrifugation and parasite material was washed twice in phosphate buffered saline/glucose (PBSG) buffer of pH 8.0, ionic strength 0.145 containing 1% glucose, Schmatz et al., J. Protozool, 31:181-183, 1984. The parasite mixture was applied to a DE52 anion exchange column, equilibrated in PBSG, and purified sporozoites were eluted unretarded in the void volume (Schmatz et al., supra).

5' GCC CAA GAA TTC GGN 3'
ALA GLN GLU PHE GLY

The 3' terminal N constitutes the first nucleotide of the cDNA and may represent any nucleotide with the resultant amino acid always being glycine.

The preferred plasmid pJC264 is derived from the plasmid pJC220 which is in turn derived from a construct containing a portion of the *E. coli* chemotaxis gene, CheY, and the gene for rat atrial natriuretic factor (ANF). The CheY-ANF plasmid is constructed from the pLC1-28, Col EI-derived plasmid described in Matsumura et al., *J. Bacteriol* 160: 36-41 (1984). The Che operon fragment containing CheY and CheZ genes is excised from the pLC1-28 plasmid as a BamHI-HindIII fragment and subcloned into a BamHI-HindIII digested pUC13 plasmid (PL Biochemicals) to give a pUC13-CheY-CheZ plasmid. *Escherichia coli* JM105 clones transformed by pUC13-CheY-CheZ express CheY and CheZ polypeptides off the lac promoter contributed by the pUC13 vector. Davis et al., *Basic Methods In Molecular Biology*, Elsevier, New York, New York, pg. 30 (1986). The pUC13-CheY-CheZ plasmid is digested at the unique PstI site internal to the CheY coding region, see Matsumura et al., *supra*, and at the unique SmaI site in the pUC13 polylinker 3' to the inserted Che DNA. The resulting 3 kb PstI-SmaI fragment containing the pUC13 vector and the DNA encoding the N-terminal 100 residues of CheY was recombined with the 160 bp PstI-HindIII fragment of pSCN1-(rat-ANF-26) that encodes the Met-(rat-ANF-26) sequence and contains 50 bp of untranslated RAS1 sequence 3' to the termination codon for the ANF peptide. This expression vector is termed the CheY-ANF vector. The pSCN1-(rat-ANF-26) fusion plasmid is constructed from the pSCN1 plasmid which expresses the N-terminal 165 amino acids of the yeast RAS1 protein SCIN, Temeles et al., *Nature* 313: 700-703 (1985). Plasmid pSCIN is digested to completion with AccI, and the ends are filled in with *E. coli* DNA polymerase I large fragment (Klenow polymerase). A synthetic ANF gene is ligated to pSCIN and used to transform competent *E. coli* JM105 cells. The nucleotide sequence of the CheY-ANF plasmid from the EcoRI restriction site to the first HindIII restriction site prior to the CheY fragment is identical to that shown for pUC19 by Yanisch-Perron et al., *Gene* 33: 103-119 (1985).

The pJC264 expression plasmid contains a unique EcoRI site, in the same reading frame as the lambda gIII EcoRI site, which permits easy subcloning and expression of EcoRI fragments from lambda gIII expression libraries. The inclusion of a portion of the CheY gene product in the resulting fusion protein may facilitate stabilization of the protein and enhance the purification of the protein. The small size of the CheY protein compared with other fusion carriers such as β -galactosidase, permits a more favorable molar yield of the protein of interest for a given mass of fusion protein. The CheY containing plasmid pJC264 results in high expression levels of fusion proteins with the first 93 amino acids of the amino terminus being derived from the *E. coli* CheY protein and linkers. As noted above the pJC264 plasmid is derived from the CheY-ANF plasmid as shown in Figure 7. CheY-ANF is partially digested with HindIII and electrophoresed in about 0.7% Seaplaque agarose gel. Full-length linear DNA is mechanically excised, removed from the gel by melting, purified on a NACS column (BRL) and recovered by ethanol precipitation. The DNA fragment is made blunt by filling in the HindIII ends with the Klenow fragment of DNA Polymerase I (Boehringer Mannheim), phenol-extracted and ethanol precipitated. BamHI linkers phosphorylated at the 5' position are ligated to the purified DNA, and *E. coli* HB 101 is transformed directly with the ligation mix. Ampicillin-resistant transformant colonies are restriction-mapped for the BamHI linker. A colony designated pJC220 contains the BamHI linker in place of the promoter-proximal HindIII site. The plasmid now has a HindIII site at the 3' end of the CheY coding region and is therefore unique. Plasmid pJC220 is digested with HindIII and two bases of the four-base overhang are filled in with the Klenow fragment of DNA Polymerase I in the presence of dATP and dGTP. The remaining two bases of the overhang are removed with SI nuclease, leaving a blunt end. The DNA is then digested with EcoRI and filled in with the Klenow fragment of DNA Polymerase I in the presence of dATP and dTTP. The plasmid is recircularized by blunt-end ligation with T4 DNA ligase to yield pJC264, which contains a unique EcoRI site at the 3' end of the CheY coding region. The new EcoRI site is in the same reading frame as the EcoRI site of lambda gIII, permitting direct subcloning and expression, as CheY fusion proteins, of antigens identified by expression in lambda gIII libraries. The pJC264 restriction map is shown in Figure 8.

Minipreps of recombinant γ gIII bacteriophage are prepared and phage DNA is isolated. The gene insert for each antigen is removed by EcoRI digestion and fractionated from the phage arms by agarose gel electrophoresis. The genes are then inserted into the plasmid pJC264 which has been linearized at its unique EcoRI site and phosphatased to decrease the efficiency of autoligation. Ligation products are then transfected into the bacterial host, *E. coli* JM83 using standard CaCl_2 methods known in the art and the

(available from Bethesda Research Laboratories) to determine which had the ability to cleave the cDNA inserts. A mapping analysis is conducted by doing single and double digests of the insert/plasmid. DNA fragments are separated electrophoretically on about 1% agarose gels, and sized by comparison to DNA markers which are run simultaneously on the same gels. Maps are constructed of each clone by entering the fragment size data and known vector restriction sites into the Intelligenetics Restriction Map Generator program (MAP, Intelligenetics, Inc.). The derived location along the nucleotide sequence of the enzymatic cleavage sites is accurate to about the $\pm 10\%$ level. The restriction maps of the group A clones are shown in Figure 1. The SO6 gene is about 1886 nucleotides (nt) in length with restriction sites at the following base locations: 118 (ApaI), 284 (PstI), 293 (PvuII), 597 (PstI), 1283 (PstI), 1820 (HincII) and 1837 (AvaI). The SP1 gene is about 1404 nt with restriction sites at the following base locations: 213 (PstI), 889 (PstI), 1386 (HincII) and 1398 (AvaI). The SO67 gene is 822 nt in length with restriction sites at the following base locations: 108 (PstI), and 816 (HincII).

The restriction maps of the group B clones are shown in Figure 2. The SO9 gene is about 1071 nt in length with restriction sites at the following base locations: 297 (PvuII), 381 (BglI), 570 (ApaI), 750 (BglI), 789 (XhoI) and 900 (PvuII). The SO24 gene is about 1108 nt in length with restriction sites at the following base locations: 243 (PvuII), 278 (BglI), 482 (ApaI), 646 (BglI), 694 (ApaI), 718 (XhoI), 743 (AvaI), 845 (PvuII) and 982 (ApaI). The SO7 gene is about 980 nt in length with restriction sites at the following base locations: 115 (PvuII), 150 (BglI), 361 (ApaI), 518 (BglI), 561 (XhoI), 564 (AvaI), 717 (PvuII) and 861 (ApaI). The SO1 gene is about 337 nt in length with restriction sites at the following base locations: 75 (ApaI), 236 (BglI), 261 (XhoI) and 275 (AvaI).

The restriction maps of the group C clones are shown in Figure 3. The SP54 gene is about 687 nt in length with restriction sites at the following base locations: 187 (AvaI), 273 (ApaI), 559 (PstI) and 627 (HindIII). The SP59 gene is about 1017 nt in length with restriction sites at the following base locations: 222 (AvaI), 250 (AvaI), 500 (AvaI), 603 (ApaI), 682 (ApaI), 889 (PstI) and 947 (HindIII).

The restriction map of a group H clone is shown in Figure 4. The SO311 gene is about 684 nt in length with restriction sites at the following base locations: 154 (HincII), 262 (BglI) and 400 (PvuII). The SO227 gene is 631 nt in length with restriction sites at the following base locations: 257 (HincII), 369 (BglI) and 537 (PvuII). The SO231 gene is 632 nt in length with restriction sites at the following base locations: 255 (HincII), 382 (BglI) and 514 (PvuII).

The restriction map of a Group F clone is shown in Figure 5. The SO216 gene is about 487 nt in length with restriction sites at the following base locations: 49 (HpaII), 97 (HhaI), 132 (KpnI), 139 (Sau3A), 176 (AluI), 200 (Sau3A), 228 (AluI), 237 (HaeIII), 296 (TaqI), 335 (HinfI), 341 (TaqI), 402 (HindIII), 404 (AluI), 415 (HhaI), 432 (TaqI), 435 (XhoI), 435 (Sau3A), 455 (HinfI) and 477 (AluI). The first eight nts and the last eight nts represent the linker nts and are not part of the *E. tenella* Group F gene.

Production of recombinant immunogenic coccidial proteins, recombinant fusion proteins and recombinant CheY fusion proteins, with recombinant CheY fusion proteins being preferred, is accomplished by overnight culturing, in 2 x YT medium containing ampicillin, of selected recombinant bacteria isolated from a single colony. The overnight culture is used to inoculate about 500 ml of 2 x YT plus ampicillin. The culture is grown at about 37°C with aeration until the mid-logarithmic phase of growth is reached, at which time IPTG is added to a final concentration of about 100 μ M. The cells are incubated for about another 3 to 4 hours, chilled on ice and collected by centrifugation. The cells are washed, collected by centrifugation and resuspended in about 10 ml of Buffer A which consists of about 30 mM Tris-HCl, about pH 8.0, about 5.0 mM EDTA and about 1 mM phenylmethylsulfonylfluoride. The cell suspension is sonicated while maintained in an ice bath in three minute bursts using a Branson cell disrupter Model 350. The sonicate is clarified by centrifugation at about 27,000 x g for about 45 minutes at about 4°C. This constitutes the first supernatant fluid. The pellet (P₁) is washed in about 10 ml of buffer A containing 0.1% w/v Triton X-100 for about 30 minutes in an ice-bath and recentrifuged. The supernatant fluid is collected and designated the second supernatant. The pellet (P₂) is washed twice in the same buffer, buffer A. The washes are discarded. The washed pellet, P₂ is then resuspended in about 1.0 ml of about 6 M guanadine-HCl containing about 100 mM dithiothreitol and the suspension incubated at about 50°C (for about 2 hours). The suspension is diluted to 10 ml with about 7 M urea and is clarified by centrifugation at about 27,000 x g for about 45 minutes at about 4°C with the supernatant fluid constituting the third supernatant. Due to differences in solubility of the various fusion proteins, some are found in the first supernatant, some in the second supernatant and some are found in the third supernatant. For example, a representative clone protein from immunogen group A, SO6-CheY, was found in the first, second and third supernatants. Representative proteins from clones of group B (SO7), C (SP54), H (SO311) and F (SO216) were found in the third supernatant. Both the SO7-CheY and SP54-CheY fusion proteins were unretarded by chromatography on hydroxyapatite. The SO311-CheY fusion protein bound to hydroxyapatite, and could be eluted with 160 mM

phosphate buffer. The SO6-CheY fusion protein from the third supernatant fluid was further purified by Trisacryl M-DEAE chromatography.

Representative *Eimeria* immunogen clones are assayed to determine the nucleotide sequence of each specific gene by one or more of three standard techniques. In some cases the nucleotide sequence of the cDNAs is determined using the chemical degradation method of Maxam and Gilbert, *Methods in Enzymology*, 65 (part 1): 497-559 (1980). More routinely, the nucleotide sequence is determined by the dideoxy chain termination technique, using denatured plasmid templates (plasmid pUC18, containing assorted subsequences of the *Eimeria* cDNAs) as described by Hattori and Sakaki, *Analyt. Biochem.*, 152:232-238 (1986). Finally, some nucleotide sequences are determined by subcloning the cDNA insert, or portions of it, into bacteriophage mp18 and sequencing secreted single-standard recombinant phage templates using the standard dideoxy chain-termination sequencing methodology of Messing, *Methods in Enzymology* 101:20-78 (1983). In addition to AMV reverse transcriptase and the Klenow fragment of DNA polymerase I, a modified T7 DNA polymerase has been employed, see Tabor and Richardson, *Proc. Nat. Acad. Sci. USA* 84:4767-4771 (1987).

The amino acid sequence is deduced from the determined nucleotide sequence by combining the following information. Each of the cDNAs in the phage expression vector γ gt11 was identified using polyclonal antisera when expressed as a fusion protein with β -galactosidase. The fusion junction between β -galactosidase and the immunogen consists of a Glu residue linking the carboxy-terminus of β -galactosidase with a Phe residue at the N-terminus of the immunogen (within the linker region). The EcoRI restriction enzyme cleaves between the first and second nucleotide of the Glu codon when reading from the 5' to 3'. This junction (and reading frame, cloning site), at the EcoRI cleavage site, is regenerated in each subsequent cloning event involving the entire cDNA irrespective of the subcloning vector, pUC18, mp18 or pJC264. Consequently, the reading frame can be unequivocally identified and the nucleotide sequence translated once the orientation of the insert in these three vectors is established. The orientation of the cDNA insert in plasmid, pUC18 and pJC264, or phage, mp18, vectors is accomplished by restriction enzyme mapping, known in the art. Once asymmetric restriction enzyme recognition sequences are identified within the cDNA insert, insert orientation and transcriptional orientation can be unequivocally assigned when the recognition sequences are similarly predicted by the nucleotide sequence. All amino acid sequences depicted herein read from the amino terminus to the carboxyl terminus.

Group A clone nucleotide sequences and the resulting Group A immunogen amino acid sequences are exemplified by the representative clone SO67. This clone is entirely contained within the SO6 clone. Of the approximately 870 nucleotides in this clone the first 162 nucleotides starting at the 5' end have been sequenced. The transcriptional orientation and therefore the correct reading frame can be deduced unambiguously based upon the location in the nucleotide sequence of restriction enzyme recognition sequences which are predicted by restriction enzyme mapping of the CheY-SO67 recombinant plasmid. The nucleotide sequence and the resulting 53 amino acid sequence is shown in Table 6. An additional 221 nucleotide sequence, see Table 7, has been obtained from the 3' end of the clone but the reading frame has not been deduced.

Group B clone nucleotide sequences and the resulting Group B immunogen amino acid sequences are exemplified by the representative clone SO7. All 957 nucleotides in this clone have been sequenced. The reading frame can be deduced unambiguously by correlating the position of restriction enzyme sites asymmetrically located within the cDNA with the location of their respective recognition sequences as predicted by the nucleotide sequence analysis. The nucleotide sequence and the amino acid sequence are shown in Table 8.

Group C clone nucleotide sequences and the resulting Group C immunogen amino acid sequences are exemplified by the representative clone SP54. This clone is entirely contained within the SP59 clone. Of the approximately 700 nucleotides in this clone the first 157 nucleotides starting at the 5' end have been sequenced. The transcriptional orientation and therefore the appropriate reading frame can be unequivocally deduced by correlating restriction enzyme recognition sequences in the nucleotide sequence with their asymmetric location predicted by restriction enzyme mapping of the CheY-SP54 recombinant plasmid. The nucleotide sequence and the resulting 52 amino acid sequence is shown in Table 9.

Group H clone nucleotide sequences and the resulting Group H immunogen amino acid sequence are exemplified by the representative clone SO311. Of the approximately 650 nucleotides in this clone, the first 185 nucleotides at the 5' end have been sequenced. The transcriptional orientation and therefore the appropriate reading frame can be unequivocally deduced by correlating restriction enzyme recognition sequences in the nucleotide sequence with their asymmetric location predicted by restriction enzyme mapping. The nucleotide sequence and the resulting 61 amino acid sequence is shown in Table 10. The last 283 nucleotides at the 3' end have been sequenced but the reading frame has not been deduced (see

Table 11).

The molecular weights of the primary translation products encoded for by the cDNAs described above are determined by *in vitro* translation of the appropriate mRNA populations. *In vitro* translation of mRNA extracted from unsporulated oocysts, sporulating oocysts and sporozoites was performed using the rabbit reticulocyte cell free translation system, with either ^{35}S -methionine or ^3H -leucine as the incorporated indicator isotope. Specific *in vitro* translation products were immunoprecipitated using monospecific antibodies, as described in Example 6. The protocol for *in vitro* translation was as described in the technical bulletin from Promega Biotec (according to manufacturer's instructions) and for immunoprecipitation as in Taylor et al., Mol. Biochem. Parasitol. 10:305-318 (1983). The *in vitro* translation product immunoprecipitated by antibody specific for the Group A antigen, exemplified by clones SO6 and SO67 has a molecular weight of about 24 kiloDaltons (kD). The *in vitro* translation product immunoprecipitated by antibody specific for the Group B antigen, exemplified by clone SO7 has a molecular weight of about 28 kD while the minor immunogens have molecular weights of about 170, 24, 22, 16 and 12 kD. The additional minor specifically immunoprecipitable *in vitro* translation products are detectable when ^3H -leucine is used as the labelled precursor amino acid. The 170 and 22 kD minor immunogens are also detectable with ^{35}S -methionine. The major 28 kD immunogen is detectable only when ^3H -leucine is used as the precursor amino acid. The *in vitro* translation product immunoprecipitated by antibody specific for the Group C antigen, exemplified by clones SP54 and SP59 has not been determined. The *in vitro* translation product immunoprecipitated by antibody specific for the Group H antigen, exemplified by clone SO311 has a molecular weight of about 28 kD while the minor immunogens have molecular weights of 48, 38, 33, 16, 13, 12 and 10 kD. The additional minor specifically immunoprecipitable *in vitro* translation products are detectable when ^{35}S -methionine is used as the labelled precursor amino acid. The major 28 kD immunogen is detectable when both ^{35}S -methionine and ^3H -leucine are used.

The specific mRNAs extracted from sporulated oocysts and/or sporozoites of *E. tenella* were sized by Northern blot analysis according to the method of Maniatis et al., Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, pg. 202 (1982) and the method described in Transfer and Immobilization of Nucleic Acids to S & S Solid supports, published by Schleicher and Schuell, Inc., pgs. 16-19 (1987). The mRNA encoding the A immunogen, exemplified by clones SO6 and SO67, is 2.15 ± 0.13 kilobases (kB) in length. The mRNA encoding the B immunogen, exemplified by clones SO7, is 1.23 ± 0.22 kB in length. The mRNA encoding the C immunogen, exemplified by clones SP54 and SP59, is 1.12 ± 0.08 kB in length. The mRNA encoding the H immunogen, exemplified by clone SO311, is 0.98 ± 0.07 kB in length.

Native immunogens, B and C are isolated from *E. tenella* by either gel filtration and identification with specific anti-CheY immunogen antibody or immuno-affinity chromatography using specific anti-CheY immunogen antibody. *Eimeria tenella* sporulated oocysts, about 1×10^9 , are sonicated in a buffer, preferably phosphate buffered saline, containing about 0.1 mM PMSF for about 10 minutes, in about 2.5 minute bursts in an ice bath. The disrupted sporulated oocysts are collected by centrifugation at $27,000 \times g$ for 30 minutes at 4°C . The pellet is washed about 3 times with about 40 ml of PBS containing about 0.1 mM PMSF and recovered by centrifugation as described above. The washed pellet is resuspended in about 60 ml of about 5M guanidine-HCl/ about 0.5 M Tris-HCl, pH about 8.6, and about 400 mg dithiothreitol. Reduction was allowed to proceed for about 3 hours at 20°C with mild agitation. Reduced and solubilized immunogen is obtained by centrifugation and collection of the supernatant fluid. The immunogen is concentrated to about 20 ml, preferably by ultrafiltration, and carboxymethylated by the addition of iodoacetic acid, about 400 mg. The pH is adjusted to about 8.6 by the addition of 3 M Tris base and the reaction allowed to continue for about 60 minutes at about 20°C in the dark. The guanidine-HCl is removed by dialysis against about 0.05 M NH_4HCO_3 , about 0.1 mM PMSF and about 0.02% sodium azide for about 48 hours. All insoluble material is removed by centrifugation. The supernatant fluid is concentrated by ultrafiltration and separated by gel filtration chromatography. The sample is applied to a column of Sephacryl S-200, about 87×2.5 cm, equilibrated in about 0.05 M NH_4HCO_3 , about 0.1% Zwittergent 3-12 and about 0.02% sodium azide. Fractions, about 4.5 ml, are collected at a flow rate of about 25 ml per hour and monitored at about 280 nm. The presence of *E. tenella* immunogen is determined by Western blotting, with rabbit anti-sporozoite antiserum and with antibody raised against the specific *E. tenella* recombinant fusion immunogens. The native immunogens are able to protect chickens against a coccidiosis infection.

Native *E. tenella* immunogens, A, B, C, H and F are isolated and purified from sporulated oocysts by immunoaffinity chromatography using antibody raised against the specific fusion immunogens. Affinity columns are prepared using preimmune serum and the specific fusion immunogen serum. Immunoglobulin G (IgG) fractions are prepared by the method of Corthier et al., J. Immunol. Meth. 66: 75-79 (1984) or by the carbonyldiimidazole method of Hearn et al., J. Biol. Chem. 254:2572-2574 (1979). About 15 mg of IgG is

coupled to 0.5 gm of Sepharose-Protein A (Sigma) using the method of Schneidert et al., J. Biol. Chem. 257:10766-10769 (1982). Approximately 5 mg of the reduced, carboxymethylated extract of *E. tenella* sporulated oocysts, prepared as described above, in about 0.1 M borate buffer, pH 8.1, about 0.5 M NaCl, about 0.02% sodium azide, and about 0.1 mM PMSF, is applied to the prebleed column equilibrated in the same buffer. The prebleed column was washed with 3 ml of column buffer and the combined column flow-through and washes are applied to the anti-*E. tenella* fusion immunogen column equilibrated in the same buffer. The column is washed with about 10 ml of column buffer and the native immunogen is eluted with about 3 M sodium thiocyanate. The individual native immunogens are able to protect chickens against a coccidiosis infection.

Immunogens from other species of *Eimeria* which share at least one antigen determinant or epitope are identified and isolated using antibody specific for *E. tenella* Group B immunogen. Other species which may share one or more common immunogens may include *E. acervulina*, *E. mivati*, *E. mitis*, *E. preacox*, *E. hagani*, *E. necatrix*, *E. maxima* and *E. burnetti*. The antibody is prepared as described above and may be either polyclonal or monoclonal. Immunogens used to product the antibody include native Group B immunogen or recombinant proteins expressed from any of the Group B clones, with the S07 clone immunogen being preferred. The recombinant immunogens may be either the individual protein or a fusion protein, with the S07-CheY fusion protein immunogen being preferred.

Immunogens associated with the various *Eimeria* species which share one or more epitopes with Group B *E. tenella* immunogen are identified by immunoblotting of immunogens prepared from sporulated oocysts and sporozoites of each individual species. The sporulated oocysts and sporozoites are physically disrupted, the proteins separated by SDS-polyacrylamide gradient gel electrophoresis and transferred to nitrocellulose. The transferred proteins are reacted with the anti-S07-CheY antibody and bound antibody is detected with ¹²⁵I-protein A.

Native Group B immunogens are isolated from *Eimeria* species sporulated oocysts by sonication, reduction and carboxymethylation. The reduced and carboxymethylated proteins may be pre-purified by size exclusion chromatography. Immunoaffinity matrices containing anti-Group B antibody are prepared using the technique of Bethel et al., J. Biol. Chem., 254: 2572-2574(1979) as described above. The Group B immunogens isolated from *Eimeria* species other than *E. tenella* are able to protect chickens against a coccidiosis infection.

Molecular weights and isoelectric points of *Eimeria* immunogens are also determined. Molecular weights are determined by analytical sodium dodecyl sulfate (SDS) polyacrylamide gel electrophoresis (PAGE) of samples prepared from sporulated oocysts and/or sporozoites of *E. tenella*, followed by transfer to nitrocellulose and immunodetection by Western Blotting as described above. Appropriate molecular weight controls are included. Isoelectric points were determined by Western Blotting of two dimensional gels run according to the procedure of O'Farrell, J. Biol. Chem. 250:4007-4021 (1975). Antibodies for both procedures are prepared as stated above. Immunogen A separated as a single band with a molecular weight of 24 kiloDaltons (kD). The predominant B immunogen is characterized as a diffuse doublet of 27-28 kD on SDS-PAGE with the minor immunogens appearing as faint bands suggesting some sharing of antigenic determinants within *E. tenella*. The minor bands have molecular weight of 22, 19, 18, 14, 12, 9, and 6 kD. The 27-28 doublet produces multiple spots on isoelectric focusing, in the range between pH 5.1 and 6 kD. The pIs of the faint additional bands detected by Western blotting were not determined. Immunogen C also migrates as a doublet with molecular weights of 21-22 kD. Immunogen H separates as two distinct major proteins with molecular weights of 28 and 18 kD and seven minor proteins with molecular weights of 27, 24, 23, 17, 14, 12, and 9 kDs. The Group F immunogen has a molecular weight of about 26-29 kD. The isoelectric points of immunogens A is 3.65 and H is 6.65. The isoelectric points of C and F have not been determined.

Poultry are administered an immunizing dosage of one of more of the recombinant derived *Eimeria tenella* immunogens described above. Immunogen administration to chickens may be by oral or parenteral routes or chicken embryos may be inoculated through the egg shell. Administration of immunogen by any of these routes may include an immunogen or immunogens given alone or as a solution or suspension with a physiologically acceptable medium. Such physiologically acceptable media include, but are not limited to, physiological saline, phosphate buffered saline, phosphate buffered saline glucose, buffered saline and the like. Parenteral administration includes inter alia, intramuscular, intraperitoneal, subcutaneous and intravenous injection or delivery of the *E. tenella* immunogens. Orally administered immunogens can be in the form of an aqueous solution or suspension. A suspension may include the immunogen in a gel composed of, for example, gelatins or alginates. Orally administered immunogens may also be included in the feed. Embryonated eggs are immunized by the injection of an immunogenic dose of one or more of the *Eimeria* immunogens. The immunogens for intramuscular and subcutaneous vaccination may be given along with an

transformants are selected on ampicillin plates. Ampicillin resistant colonies are grown on an analytical scale to score for the presence of an insert, score for orientation of the foreign DNA with respect to the bacterial promoter and score for expression of bacterial fusion proteins by Western blot analysis, using polyclonal antisera raised against *E. tenella* immunogens.

DNA inserts are isolated from phage clones representative of the various immunogen groups identified above and are also sub-cloned into the puc18 plasmid-vector as described above for the CheY vector, pJC264. Restriction endonuclease maps of members of each group are prepared. The restriction endonucleases included, but are not limited to, the following:

AluI	HindIII	Sall
ApaI	HincII	Sau3a
AvaI	HinfI	SstI
Avall	HpaII	SstII
BamHI	KpnI	TaqI
BglI	NcoI	XbaI
Clal	PstI	XhoI
HaeIII	PvuI	XhoII
HhaI	PvuII	

all of which are available commercially. The following table contains the groups, clone designation within each group and the restriction endonucleases which are unable to cut within the clone insert.

TABLE 3

RESTRICTION ENDONUCLEASE SITES ABSENT FROM DESIGNATED CLONES		
Group	Clone Designation	Restriction Endonucleases
A	SO6' SPI SO67	BamHI, HindIII, KpnI, NcoI, Aval, Clal, XhoI, Sall, SstI, SstII, XbaI, BglI,
B	SO9 SO24 SO7' SO1'	BamHI, HincII, KpnI, NcoI, Clal, Sall, SstI, XbaI
C	SP54 SP59	BamHI, KpnI, HincII, NcoI, Clal, PvuII, XhoI, Sall, SstI, SstII, XbaI, BglI
H	SO311 SO227 SO231	BamHI, HindIII, KpnI, Avall, ApaI, NcoI, Aval, Clal, PstI, XhoI, Sall, SstII, XbaI
F	SO216	ApaI, Aval, Avall, BamHI, BglI, Clal, HincII, NcoI, PstI, PvuII, Sall, SstI, SstII, XbaI, XhoI

Some restriction endonucleases are capable of cleaving one or more clones within a group but not all clones. In the B group, additional restriction endonucleases which cleave at least one of the four clones include Aval, PstI, SstII. These sites have not been mapped. In the H group, the restriction endonuclease SstI does cleave within all three of the clones, but the site has not yet been mapped.

The above information is determined by growing the pUC 18 recombinant plasmids as mini-preparations in LB broth and isolating the DNA using the alkaline lysis method described below. The DNA is resuspended in digestion buffer such as TE buffer which contains, about 10 mM Tris-HCl (about pH 8.0), about 1 mM EDTA (about pH 8.0), containing DNase-free pancreatic RNase, about 20 µg/ml and mixed on a Vortex mixer briefly. The DNA samples are then digested with a variety of restriction endonucleases

mM NaCl/about 100 mM Tris-HCl, about pH 7.5, about 5 mM MgCl₂. The reaction products are adjusted to about 0.3 M sodium acetate by the addition of about one-tenth volume of a 3M (pH 5.6) stock solution, precipitated with ethanol, chilled and collected by centrifugation. After suspending the pellet in TE, the DNA is electrophoresed in agarose containing ethidium bromide to resolve the insert from the phage arms.

Fractionation of the inserts is verified by visualization under ultraviolet light. The inserts are electrophoresed onto NA-45 (Schleicher & Schuell) membranes and then eluted from the membranes. Insoluble particles are removed by centrifugation and the soluble material is extracted with phenol, phenol/chloroform/isoamyl alcohol and chloroform/isoamyl alcohol. The DNA is precipitated with sodium acetate/ethanol, washed with ethanol and air dried. An aliquot of each DNA is analyzed on an analytical agarose gel for confirmation.

Expression of the genes coding for the protective coccidial immunogens is accomplished in a number of different host cells with a variety of promoter-expression systems. The host cells include bacteria, yeast, insect, and mammalian cells. The antigens may also be expressed in a number of virus systems. Although the genes can be expressed in numerous procaryotic cells and various eucaryotic cells the most preferred host cell is Escherichia coli. The expression vectors which can be used for the expression of the protective immunogens include, but are not limited to, pBR322, pPLa2311, pKC30, ptac12, γ gt11, pAS1, pLC24, pSB226, pRIT2T and SV40 with a CheY-pUC derived vector designated pJC264 being preferred. It is desired and intended that there be included in this invention, the use of Eimeria tenella immunogens, which are native proteins or fragments thereof, recombinant proteins or fragments thereof, or fusion proteins linked to other proteins which may or may not enhance the Eimeria peptides immunogenicity. The fusion immunogens may be designed in such a manner that the immunogenic expression protein contains an additional polypeptide portion encoded for by the expression-plasmid or an additional peptide portion that has been added to the gene by the inclusion of an additional DNA base sequence. The pJC264 plasmid is designed to include the expression of an 88 amino acid portion of the E. coli CheY protein operably attached to 5 linker amino acids linked or fused to the various Eimeria tenella peptides. Operably attached refers to an appropriate sequential arrangement of nucleotide segments, linkers, or genes such that the desired protein will be produced by cells containing an expression vector containing the operably attached genes, segments or linkers. The nucleotide sequence of the CheY gene and the amino acid sequence produced from the gene are shown in the following table.

EXAMPLE 4Preparation of Genomic DNA from *Eimeria tenella* Sporozoites

Purified *Eimeria tenella* sporozoites, from Example 1 were suspended in TE medium (10 mM Tris-HCl, pH 7.5, 0.1 mM EDTA) at a concentration of 1.5×10^8 sporozoites per ml. The dilute suspension of sporozoites was then adjusted to 0.5% in SDS (from a 20% SDS stock solution), and 15 mM in EDTA (from a 0.5 M-pH 8.0 stock solution) which resulted in both plasma and nuclear membrane lysis. The release of genomic DNA following nuclear lysis is marked by an obvious increase in the viscosity of the solution. To aid in solubilization, the solution was gently rocked at 50°C on a platform for 30-60 minutes, and then digested for 3 hours at 50°C with Proteinase K at a concentration of 100 ug per ml. Genomic DNA was purified by two extractions with phenol, two extractions with a mixture of phenol, chloroform and isoamyl alcohol (25:24:1), two extractions with chloroform and isoamyl alcohol (24:1), and two successive precipitations with sodium acetate/ethanol as described in Example 8. The nucleic acid pellet was washed twice with 70% ethanol and suspended in TE at an approximate concentration of 5×10^8 sporozoite equivalents per ml. The RNA component of the nucleic acid was selectively removed by digestion with heat inactivated RNase A at a concentration of 50 ug per ml for 60 minutes at 37°C. The RNase A and other residual proteins were removed by a secondary digestion with Proteinase K in 0.5% SDS and 15 mM EDTA for 3 hours at 50°C as described above. Genomic DNA was then successively extracted with organic solvents, precipitated twice with ethanol, and then washed twice with 70% ethanol. The genomic DNA pellet was suspended in TE at a concentration of $2-3 \times 10^9$ sporozoite equivalents per ml and quantitated by absorbance at 260 nm. Undigested genomic DNA was then fractionated on an analytical gel to confirm (i) the spectrophotometric-derived concentration, (ii) the lack of residual RNA, and (iii) its high molecular weight integrity.

EXAMPLE 5Construction of cDNA Expression Libraries

E. tenella oocysts, sporulated for seven hours, and sporozoites were prepared as previously described (Schmatz et al, supra; Wang & Stotish, J. Protozool. 22: 438-448, 1975). Total RNA was isolated from each stage either immediately after isolation (i.e. the sporozoites) or from cell pellets frozen in liquid nitrogen and stored at -80°C (i.e., the 7 hour sporulating oocysts) by the method of Chirgwin et al., (Biochem. 18: 5294-5299, 1979). Due to the presence of the cell wall, oocyst samples were resuspended in 4 volumes of 4 M guanidinium thiocyanate solution (volumes of solution relative to volume of cell pellet) and were sonicated for a total of 30 minutes at 20 W, 50% cycle with a Branson sonifier (Heat System Ultrasonics). Sporozoites were lysed upon the addition of the guanidinium thiocyanate stock solution (4 M guanidinium thiocyanate, 0.5% N-lauroylsarcosine, 25 mM sodium citrate, pH 7.0, and 0.1 M 2-mercaptoethanol); therefore sonication was unnecessary. The lysed cells were then centrifuged at 8,000 rpm for 10 minutes in a Beckmann JS-13 rotor at 10°C to sediment particulate cellular debris. The supernatants were decanted into a clean flask and mixed with 0.025 volumes of 1 M acetic acid and 0.75 volumes of absolute ethanol. The flask was shaken thoroughly and left to stand overnight at -20°C to precipitate the nucleic acids. The next day, the RNA was collected by centrifugation in a Beckmann JS-13 rotor at 8000 rpm for 10 minutes at 10°C. The tubes were drained and the cell pellet was resuspended in 0.5 volumes of buffered guanidine hydrochloride stock solution (7.5 M guanidine hydrochloride, 0.025 M sodium citrate, pH 7.0, and 5 mM dithiothreitol). The volume of the guanidine hydrochloride stock solution is relative to the volume of the guanidinium thiocyanate solution previously used. The RNA was precipitated by adding 0.025 volumes of 1 M acetic acid and 0.5 volumes of absolute ethanol. The solution was kept overnight at -20°C and the RNA was collected once again by centrifugation. The guanidine hydrochloride precipitation was repeated, using half the volume of the guanidine hydrochloride stock solution used in the previous precipitation. The reprecipitated RNA was washed in 95% ethanol, dried, and resuspended in sterile water. This material was centrifuged for 30 minutes at 10,000 rpm (Beckmann JS-13 rotor) at 10°C. The supernatant fluids were saved and the pellets were resuspended in sterile water. The centrifugation step was repeated. The supernatant fluids were combined, mixed with 0.1 volume of 2 M potassium acetate, pH 5, and 2 volumes of absolute ethanol, and were left to precipitate overnight at -20°C. The RNA pellets were collected by centrifugation at 10,000 rpm (Beckmann JS-13 rotor) for 30 minutes, dried, and resuspended in sterile water. The concentration of the RNA was determined by spectrophotometry.

Polyadenylated RNA was selected by oligo (dT)-cellulose chromatography (Aviv & Leder, Proc. Nat. Acad. Sci. USA 69: 1408-1412, 1972). To make a 1 ml column, 0.3 g of oligo (dT)-cellulose (Bethesda Research Laboratories, BRL) was resuspended in elution buffer (10 mM Tris-HCl, pH 7.5) and poured into a Pasteur pipette. Before use, the column was washed with 10 bed volumes of binding buffer (0.5 M lithium chloride, 0.5% sodium dodecyl sulfate, 10 mM Tris-HCl, pH 7.5, and 1 mM ethylenediamine tetraacetic acid).

The RNA (0.5 mg), dissolved in sterile water, was heated at 68°C for 5 minutes and cooled to room temperature on ice. An equal volume of 2X binding buffer was added, mixed thoroughly, and the sample was applied to the column. After washing the column with 50 mls of binding buffer, the poly(A⁺)-RNA was eluted with 10 mls of elution buffer. Ten, 1 ml fractions were collected and the concentration of RNA in each was determined by spectrophotometry at a wave-length of 260 nm. The fractions with the highest absorbance were pooled and RNA was precipitated by adding 0.1 volumes of 2 M potassium acetate, pH 5.0, and 2 volumes of absolute ethanol. The samples were left overnight at -20°C and the RNA was collected by centrifugation as above. After precipitation, the samples were resuspended in sterile water and the concentration of each was redetermined by spectrophotometry.

Starting with 7.5 µg of poly(A⁺)-RNA, first and second strand cDNA reactions were performed as described by Gubler and Hoffman (Gene 25:263-269, 1983). Synthesis of the first strand of the cDNA was carried out in a reaction volume of 40 µl containing 50 mM Tris-HCl, pH 8.3, 10 mM MgCl₂, 10 mM DTT, 4 mM Na-pyrophosphate, 1.25 mM dGTP, 1.25 mM dATP, 1.25 mM TTP, 0.5 mM dCTP, 15 µCi of [α -³²P] dCTP (3000 Ci/mmol), 100 µg/ml of oligo (dT₁₂₋₁₈), 3000 units AMV reverse transcriptase/ml (Beard, Life Sciences, St. Petersburg, Fl) for 30 minutes at 42°C. The products were extracted with phenol/chloroform and precipitated with absolute ethanol out of 2 M NH₄-acetate, Okayama & Berg, Mol. Cell Biol. 2: 161-170, 1982. The pellets were washed with 70% ethanol, dried, and resuspended in 40 µl of sterile water.

For second strand synthesis, 500 ng of single-stranded cDNA (i.e. 1 µg of the cDNA/mRNA hybrid) was resuspended in 100 µl of 20 mM TRIS-HCl, pH 7.5, 5 mM MgCl₂, 10 mM (NH₄)₂SO₄, 100 mM KCl, 0.15 mM β -NAD, 50 µg per ml BSA, 40 µM each of dATP, dGTP, dCTP and dTTP, 8.5 units/ml of E. coli RNase H (Pharmacia P-L Biochemicals, Inc.) and 230 units per ml E. coli DNA polymerase I (Pharmacia P-L Biochemicals, Inc.). Incubations were sequentially carried out at 12°C for 60 minutes and at 22°C for 60 minutes. EDTA was added to 20 mM to stop the reaction and the products were extracted twice with phenol/chloroform. The double stranded cDNA was precipitated with 2 volumes of absolute ethanol from 2 M NH₄-acetate as previously described.

The cDNA (50 ng-1 µg) was then methylated in a 20 µl volume of IX EcoRI methylase buffer containing 50 mM Tris-HCl, pH 7.5, 1 mM EDTA, 5 mM DTT, and 10 µM S-adenosylmethionine. The reaction was carried out at 20°C for 20 minutes after the addition of 20 U of EcoRI methylase (New England Biolabs). To terminate the reaction, the enzyme was heat inactivated for 15 minutes at 70°C. The samples were cooled on ice and the cDNA was blunt-ended as follows. To the tube containing 21 µl of EcoRI-methylated cDNA, 2.5 µl of 0.1 M MgCl₂, 2.5 µl of 0.2 mM d(A, C, G, T)TP and 5 units of T4 DNA polymerase (BRL) were added. The reaction was carried out at 20-22°C for 10 minutes and terminated with the addition of EDTA to a final concentration of 15 mM. The reaction products were extracted twice with phenol/chloroform and precipitated with ethanol as above.

The pellets from the previous reactions were resuspended in 4.5 µl of 100 µg/ml kinased EcoRI dextranucleotide linkers (BRL) in buffer containing 70 mM Tris-HCl, pH 7.6, 10 mM MgCl₂, 5 mM DTT, and 1 mM ATP. T4 DNA ligase (New England Biolabs, 200 U/0.5 µl) was added and the reaction mixture was incubated overnight at 12°C. The linker ligated cDNA's were then digested to completion with EcoRI (BRL). To the 5.5 µl overnight incubation, 5 µl of EcoRI correcting buffer (50 mM Tris-HCl, pH 7.5, 10 mM MgSO₄, 200 mM NaCl) was added. The mixture was heated for 10 minutes at 70°C to inactivate the ligase. The volume of the reaction mixture was increased two-fold (to 20 µl) with 100 mM Tris-HCl, pH 7.5, 50 mM NaCl, and 10 mM MgCl₂ and 2 µl of EcoRI restriction endonuclease (16 units/ul) was added. The digest was allowed to proceed for one hour at 37°C after which the enzyme was heat inactivated for 20 minutes at 65°C. The products were precipitated as above.

To remove the digested linkers from the reaction mixture, the cDNA was further purified on an Elutip-d column (Schleicher and Schuell). Finally, the cDNA (300 ng) was ligated into 7.5 µg of commercially purchased EcoRI-digested, alkaline phosphatase treated γ tII vector DNA (Promega Biotec). The vector-to-donor molar ratio in the ligation mixture was 1:1, and the final concentration of DNA was approximately 200 µg/ml. The ligation reaction was carried out in 10 mM Tris-HCl, pH 7.5, 10 mM MgCl₂. To anneal the cohesive ends of the γ vector, the mixture was first incubated at 42°C for 15 minutes. It was then supplemented with 1 mM ATP, 10 mM DTT, and 40,000 units/ml of T4 DNA ligase (New England Biolabs). The reaction was incubated overnight at 14°C.

The γ vector hybrids were packaged in vitro with commercially available packaging extracts according to the manufacturer's instruction (Amersham). Small aliquots of the packaged phage were transduced into *Escherichia coli* host strain Y1088 (Huynh et al., In "DNA cloning: A practical approach", Volume I, Glover, D. ed., IRL Press, Oxford, pp 49-78, 1985) and these were plated on LB plates using 2.5 ml of LB (10 g per L Bactotryptone, 5 g per L Bacto-yeast extract, 10 g per L NaCl, pH 7.5) soft agar containing 600 $\mu\text{g ml}^{-1}$ X-gal and 16 mM IPTG. Two cDNA libraries, each consisting of approximately 1×10^7 independent recombinant phage clones were generated. The nonrecombinant background, as determined by growth on X-gal/IPTG plates, was estimated to be 13%.

EXAMPLE 6

Screening of γgtII cDNA Libraries

The screening of the cDNA libraries from Example 5 with either anti-Fraction V antibody or anti-sporozoite antibody, from Example 2, was done essentially as described by Huynh et al., supra. Packaged phage from the unamplified cDNA library were transduced into *E. coli* strain Y1090 and plated on 150 mm plates at a density of $0.5\text{--}1.0 \times 10^5$ plaque forming units (pfu) per plate. The plates were incubated at 42°C for 3.5 hours, overlaid with dry nitrocellulose filters presoaked in 10 mM IPTG, and incubated overnight at 37°C . The filters were removed, blocked for 1 hour with 20% fetal calf serum in Tris buffered saline (TBS; 50 mM Tris-HCl/150 mM NaCl, pH 8.0) containing 0.05% Tween 20 (TBST), and were then incubated with the appropriate antibody for an equivalent length of time. Antibody binding sites were detected with [^{125}I] labeled protein A. Positive plaques were picked, replated, and rescreened until each clone was shown to be plaque pure.

For cross-screening experiments, 1 μl of phage lysate from each plaque purified clone was spotted on a lawn of *E. coli* Y1090 cells. Recombinant fusion proteins were induced, transferred to nitrocellulose, and immunoblotted as described below. Screening and cross-screening with the various antisera revealed the five groups of clones in Table 1. All of the antisera used for immunoblotting were exhaustively preabsorbed with a concentrated lysate of γgtII lysogen BNN93. After preabsorption, they were diluted 1:100 in TBST and stored at 4°C until required.

Monospecific antibodies to each of the recombinant phage were affinity purified from polyspecific antisera, from Example 2, by a modification of the method of Hall et al. (Nature 311: 379-382, 1984) and by immunizing rabbits as described in Example 2 with the purified recombinant *E. tenella*-CheY fusion proteins as described in Example 13. The fusion proteins included Group A, SO67-CheY; Group B, SO7-CheY, Group C, SP54-CheY; Group H, SO311-CheY; and Group F, SO216-CheY. Filter plaque lifts were prepared from purified recombinant clones as was done for screening. Approximately 2×10^5 pfu were plated per 150 mm plate to give close to semiconfluent lysis at the end of the 37°C incubation period. The nitrocellulose was then removed, blocked with 20% fetal calf serum in TBST for 4 hours, and incubated overnight with 20 ml of preabsorbed polyspecific serum (diluted 1:200 with 20% fetal calf serum in TBST containing 0.02% NaN_3). All of the incubations were done at room temperature with constant agitation. Subsequently, the filters were washed five times for 20 minutes each with 50 ml of TBST and one time with 0.15 M NaCl/0.05% Tween 20. The antibodies were eluted from each of the filters with 10 ml of 0.2 M glycine-HCl/0.15 M, NaCl/0.05% Tween 20, pH 2.8 for 30 minutes. The pH of each eluate was restored to 8.0 with Tris base and the recombinant eluted antibodies (REA's) were stored at -20°C until required.

Parasite antigens were obtained by sonicating unsporulated oocysts, sporulated oocysts, and DE-52 purified sporozoites in NET buffer (50 mM Tris-HCl, pH 7.5, 150 mM NaCl, 5 mM EDTA) with 1 mM phenylmethylsulfonylfluoride (PMSF) as a protease inhibitor as described in Example 1. Protein concentrations of each sample were determined by method of Lowry et al., supra. The yield of antigen from 3×10^5 unsporulated/sporulated oocysts was approximately 50 μg , whereas the same amount of antigen was obtained from approximately 2×10^6 sporozoites. Samples were kept at -20°C until ready for use. For blots of parasite antigens, 50 μg of each sonicated sample was mixed with an equal volume of 2x sample buffer (0.125 Tris-HCl, pH 6.8, 4% w/v SDS, 10% v/v 2-mercaptoethanol, 20% glycerol and 0.0025% bromophenol blue), boiled for 3 minutes and electrophoresed on either a 15% SDS-polyacrylamide gel or a 5-20% SDS-polyacrylamide gradient gel (Laemmli, Nature 227:680-684, 1970).

Alternatively antigens were prepared by resuspending oocysts at a concentration of 5×10^7 per ml and sporozoites at a concentration of 5×10^8 per ml in NET buffer containing a cocktail of protease inhibitors (2 mg ml^{-1} 1-10 phenanthroline, 2 mg ml^{-1} benzamidine, 0.002 mg ml^{-1} PMSF, 0.048 mg ml^{-1} Sigma soybean trypsin inhibitor, 0.048 mg ml^{-1} aprotinin, 0.02 mg ml^{-1} leupeptin). At this point the samples were mixed with an equal volume of 2x sample buffer without bromophenol blue. The samples were boiled for 3

minutes, sonicated till fully disrupted, and reboiled again for 3 minutes. Bromophenol blue was added to 0.0025% and the samples were stored at -20°C until ready for use. For immunoblotting, oocyst or sporozoite antigens were loaded and subjected to electrophoresis as stated above.

Proteins separated by SDS-PAGE were electrophoretically transferred to nitrocellulose by the technique of Towbin et al., Proc. Natl. Acad. Sci. USA, 76:4350-4354 (1979). The nitrocellulose was subsequently blocked with 20% fetal calf serum in TBST for 4 hours. After blocking, the nitrocellulose was incubated overnight at room temperature in 20 ml of antibody diluted with 20% fetal calf serum in TBST containing 0.02% NaN_3 . Polyspecific antisera were diluted 1:100 to 1:200 and monospecific recombinant eluted antisera were diluted 1:10. Following the contacting with specific antibody, the filters were washed three times for 5 minutes each with 200 ml of TBST. Bound antibody was detected with ^{125}I -protein A diluted in 20 ml of TBST to a final concentration of 2×10^5 counts per minutes ml^{-1} . Incubation with radio-labelled protein A was carried out for 1 hour at room temperature after which time the filters were again washed three times for 5 minutes with 200 ml of TBST, were air dried, and exposed to Kodak X-omat AR film.

Alternatively, the nitrocellulose was blocked with 0.5% gelatin in phosphate buffered saline, pH 7.4, for 1 hour with three 200 ml washes followed by a second blocking with 0.25% gelatin in TEN buffer, 50 mM Tris-HCl, 150 mM NaCl, 5 mM EDTA, pH 7.4, for 1 hour and washed as before. After blocking, the nitrocellulose was incubated overnight at room temperature in 20 ml of antibody diluted 1:100 to 1:200 with TEN buffer containing 0.25% gelatin and 0.05% Triton X-100. The filters were washed 5 times for 20 minutes each with 200 ml of TEN containing 0.25% gelatin. Bound antibody was detected with ^{125}I -protein A diluted in 20 ml of TEN, 0.25% gelatin, 0.05% Triton to a final concentration of 2×10^5 cpm ml^{-1} . Incubation with radiolabelled protein A was carried out for 1 hour at room temperature, after which time the filters were washed 2 times for 15 minutes with 200 ml of TEN containing 0.25% gelatin and 0.05% Triton and 4 times for 15 minutes with 200 ml of TEN. After washing, the filters were air dried and exposed to Kodak X-omat AR film.

EXAMPLE 7

Preparation of phage DNA

Recombinant and wild type γgtII phage from Example 6, were introduced as lysogens into *E. coli* host strain Y1089 (Huynh et al., supra) at a multiplicity of 10. The lysogens were streaked on to LB-plates containing $100 \mu\text{g ml}^{-1}$ ampicillin for single colony isolation and incubated overnight at 30-32°C. The growth of several colonies was checked at 32°C and 42°C. One colony was picked from a 32°C plate that did not grow at 42°C, and an overnight culture was set up in LB broth with 50 mg L^{-1} ampicillin.

The lysogenized clones were then grown from the overnight culture in 50 ml of LB broth containing $50 \mu\text{g ml}^{-1}$ ampicillin at 32°C until an O.D. 600 of 0.3 to 0.5 was reached. Phage excision and replication was induced by a temperature shift to 45°C for 20 minutes. Continued phage replication was insured by continuing to grow the cultures at 37°C for 2 to 3 hours, until sign of cell lysis was visible. If the cultures were not completely lysed, 0.1 ml of chloroform was added to each, and the cultures were agitated for an additional 10 minutes at 37°C. Under these conditions, lysis of the cells occurs after a few minutes. The cellular debris was routinely removed, at this point, by centrifugation for 5 minutes at 7,000 rpm in a Beckmann JS-13 rotor. The phage supernatant fluids were stored overnight at 4°C after adding MgSO_4 to a final concentration of 0.01 M, to stabilize the phage heads.

After bringing the phage supernatant fluids to room temperature, $50 \mu\text{l}$ of 10 mg ml^{-1} DNase I and $25 \mu\text{l}$ of 10 mg ml^{-1} RNase A were added to each sample. These were incubated for minimally one hour at 30°C, after which 1.46 gm of NaCl was added and thoroughly dissolved in each. The supernatant fluids were incubated further on ice for a minimal time of 30 minutes. The remaining cellular debris was then collected by centrifugation for 10 minutes at 10,000 rpm in a Beckmann JS-13 rotor. The supernatants were collected from each sample and in each supernatant fluid, 3.5 gm of Carbowax PEG 8000 (polyethyleneglycol 2000, Fisher Scientific Co.) was dissolved. In the presence of PEG, the phage heads were left to precipitate overnight at 4°C. The next day, the phage heads were collected by centrifugation. The supernatant fluids were centrifuged for 10 minutes at 10,000 rpm in a Beckmann JS-13 rotor maintained at 4°C. The supernatant fluids were carefully drained off and discarded. The pellets were resuspended in $250 \mu\text{l}$ of 0.1 M Tris-HCl (pH 7.9), 0.3 M NaCl, and 1 mM EDTA, after which $12.5 \mu\text{l}$ of 0.5 M EDTA was added to chelate any free Mg^{++} left behind in the sample. The phage heads were incubated in the aforementioned buffer for 10 minutes at 67°C. After the incubation, $5 \mu\text{l}$ of 10% SDS was added to each sample and the samples were mixed on a vortex mixer. Heating was used to denature the phage proteins. The SDS completes the denaturation step, and releases the DNA from the phage heads.

The DNA which has been released from the phage is then extracted twice with phenol, three times with chloroform-isoamyl alcohol (24:1), and precipitated with the addition of one-tenth volume of 3M NaOAc (pH 7.5) and two volumes of absolute ethanol. The samples were left to precipitate overnight at -20 °C. The next day, the DNA was collected by centrifugation in a microfuge for 20 minutes. The precipitated DNA was redissolved in 300 µl of 0.3 M KOAc and reprecipitated with the addition of two volumes of absolute ethanol. The samples were incubated at -80 °C for 10 minutes and the DNA was collected by centrifugation as described above. The DNA pellets were washed with 70% ethanol, dried, and resuspended in 100 µl of TE buffer (10 mM Tris-HCl (pH 7.6), 1 mM EDTA (pH 8.0)). The concentration of DNA in each sample was determined by spectrophotometry at a wavelength of 260 nM.

EXAMPLE 8

Purification of cDNA Insert From γ gtII Clones

Ten to 20 µg of γ gtII recombinant phage, from Example 7, (at a final DNA concentration of 0.2 µg/ul) was cut to completion with EcoRI (80U/µl; Boehringer Mannheim) in a reaction buffer composed of 50 mM NaCl/100 mM Tris-HCl (pH 7.5)/5 mM MgCl₂. The reaction was conducted at 37 °C for 4 hours using a 5-fold enzyme excess. Reaction products were adjusted to 0.3 M sodium acetate by the addition of one-tenth volume of a 3 M (pH 5.6) stock solution, precipitated with 2.5 volumes of ethanol, chilled for 20 minutes at -70 °C, and collected by centrifugation at 15,000 x g for 15 minutes at 4 °C. The pellet was suspended in 30 µl of TE (10 mM Tris-HCl, pH 7.5/0.1 mM EDTA) and loaded onto a preparative 1% agarose flat bed gel containing ethidium bromide. The insert was resolved from the phage arms by electrophoresis overnight (15 hr/60 mA).

Fractionation of the insert was verified by visualization under ultraviolet light. The agarose gel was sliced on both sides of the cDNA insert and pieces of NA-45 membrane (Schliecher & Schuell) were inserted into the gel, "sandwiching" the cDNA insert. The insert was then electrophoresed onto the NA-45 membrane. Upon completion, the membrane was removed from the gel, cut into small pieces and placed into an Eppendorf tube with 250 µl of a solution composed of 50 mM arginine (free base), 1 M NaCl. DNA was eluted from the membrane at 70 °C for 3 hr; the aqueous solution was removed and the elution process was repeated using a fresh 250 µl of eluant. The two eluates (totaling 500 µl) were combined and chilled to 4 °C. Insoluble particulates were collected by centrifugation for 10 minutes at 4 °C at 15,000 x g. The soluble material was then extracted twice with phenol, twice with phenol/chloroform/isoamyl alcohol (25:24:1) and twice with chloroform/isoamyl alcohol (24:1). DNA was precipitated with 0.3 M sodium acetate/EtOH (as described above), washed twice with 70% EtOH, air dried, suspended in 25 µl of TE and quantitated by absorbance at 260 nM. An aliquot of the DNA was then analyzed on an analytical agarose gel for confirmation.

EXAMPLE 9

Mapping of cDNA clones isolated from γ gtII library

DNA inserts, from Example 8, were isolated from phage clones representative of Group A (SO6', SP1, SO67), Group B (SO9, SO24, SO7', SO1'), Group C (SP54, SP59) Group H (SO311, SO227, SO231) and Group F (SO216). The phage inserts were subcloned into the plasmid vector, puc18, which is commercially available from Bethesda Research Lab. Both the isolation of inserts as well as the subcloning were done as described for the CheY vector, pJC264 in Example 12. The plasmids were grown as mini-preparations in 5 ml cultures of LB broth, and the DNA was isolated from each, using the alkaline lysis method as described in Example 12. The DNA was resuspended in 50 µl of TE buffer, 10 mM Tris-HCl (pH 8.0), 1 mM EDTA (pH 8.0), containing DNase-free pancreatic RNase (20 µg ml⁻¹) and by brief vortex mixing. The DNA samples were then digested with a variety of restriction endonucleases (commercially available from many suppliers including Bethesda Research Laboratories) in order to determine which were cutters or noncutters of the cDNA inserts. The restriction enzyme digestions were always done according to the manufacturer's recommendations. Usually five cutters were chosen for each clone, and a mapping analysis was conducted by doing single and double digests of each recombinant plasmid. The DNA fragments which were generated were separated electrophoretically on 1% agarose gels, and sized by comparison to DNA markers which were run simultaneously on the same gels. Maps were constructed of each clone by entering the fragment size data and known vector restriction sites into the Intelligenetics Restriction Map Generator program (MAP Intelligenetics, Inc.). In each case, the map which is the most compatible with all

of the data is shown in the Figures I-V.

EXAMPLE 10

5 Construction Of The CheY-ANF Plasmid

An expression plasmid for the fusion polypeptide SC1N-(rat-ANF-26) was derived from the pSCN1 plasmid. The pSCN1 plasmid is a bacterial-expression plasmid for the N-terminal 165 amino acids of the yeast RAS1 protein SCIN and is described in Temeles et al., Nature 313: 700-703 (1985). The plasmid
10 pSCIN (1 µg) was digested to completion with AccI, and the ends were filled in with E. coli DNA polymerase I large fragment (Klenow polymerase). The synthetic ANF gene was excised by digestion of pANF-I with DdeI and Hinc II. After filling out the DdeI end with Klenow polymerase, the 104 bp fragment was isolated. The ANF gene fragment was then ligated to pSC1N treated as described above and used to transform competent JM105 cells. Ampicillin-resistant colonies were screened with an appropriate
15 oligonucleotide. SDS extracts of hybridization positive colonies were electrophoresed on a 15% sodium dodecyl sulfate polyacrylamide gel (SDS-PAGE), followed by either staining with Coomassie Blue or protein blot analysis to detect the expression of the fusion protein.

The ANF gene was transferred from the pSCN1 plasmid to the pLC1-28 plasmid. Plasmid pLC1-28 is a col EI-derived plasmid that contains the entire Che operon and is described in Matsumura et al., J. Bacteriol. 160: 36-41 (1985). The Che operon fragment containing the CheY and CHEZ genes was excised
20 from pLC1-28 as a BamHI-HindIII fragment and sub-cloned into BamHI-Hind III digested pUC13(PL Biochemicals) to give pUC13-CheY-CheZ. Escherichia coli JM105 clones transformed by pUC13-CheY-CheZ expressed CheY and CheZ polypeptides off the lac promoter contributed by the pUC13 vector. To construct an expression plasmid for the CheY-(rat-ANF-26) fusion, pUC13-CheY-CheZ was digested at the
25 unique PstI site internal to the CheY coding region and at the unique SmaI site in the pUC13 polylinker 3' to the inserted Che DNA. The resulting 3 kb PstI-SmaI fragment containing the pUC13 vector and the DNA encoding the N-terminal 100 residues of CheY was recombined with the 160 bp Pst I-HindIII fragment of pSCN1-(rat-ANF-26) that encodes the Met-(rat-ANF-26) sequence and contains 50 bp of untranslated RAS1 sequence 3' to the termination codon for the ANF peptide, see Figure 6. E. coli JM105 was transformed
30 with the ligation mix containing the two fragments described above. DNA was isolated (minipreps) from ampicillin-resistant clones. The desired clones were identified as those releasing a 160 bp gene fragment upon EcoRI-Pst I digestion. These clones were shown to express ANF peptides by Western Blot analysis of total cellular protein using anti-ANF antisera.

35 EXAMPLE 11

Construction of Plasmid pJC264

The CheY-ANF plasmid from Example 10 was converted to the plasmid pJC220 which was in turn
40 modified to produce the unique pJC264 plasmid. To convert CheY-ANF to pJC220, 40 µg of CheY-ANF plasmid DNA was incubated at 37 °C with 20 units of HindIII (International Biotechnologies, Incorporated) in a final volume of 200 µl of 25 mM Tris-HCl pH 7.8, 50 mM NaCl, 10 mM MgCl₂, 1 mM dithiothreitol, and 100 µg/ml bovine serum albumin. At 15 minute intervals 50 µl aliquots were transferred to tubes containing
45 2 µl 0.5M Na-EDTA, pH 8.0 to stop the digestion. Each sample 150 ng was electrophoresed in adjacent lanes of a 0.7% (w/v) Seaplaque agarose (FMC) gel containing 89 mM TRIS, 89 mM boric acid, 2 mM EDTA (TBE) and 0.5 µg/ml ethidium bromide. The linearized plasmid was identified as that band comigrating with XhoI-digested CheY-ANF when visualized by 365 nm light. This band was excised from the gel with a razor blade from the lanes corresponding to 15, 30, 45 and 60 minutes of digestion, melted at
50 65 °C, and diluted with 10 volumes of 0.2M NaCl, 10 mM Tris-HCl pH 7.2, 1 mM EDTA, at 37 °C (Buffer A). The DNA was bound to a NACS Prepac cartridge (Bethesda Research Laboratories) BRL by gravity flow, washed with 10 ml Buffer A, and eluted with 0.5 ml Buffer D (2 M NaCl, 10 mM Tris-HCl pH 7.2, 1 mM EDTA) by gravity flow. One ml absolute ethanol was added to the column eluate. The sample was mixed and incubated on dry ice 10 minutes and centrifuged at 12,000 x g for 15 minutes at 4 °C. The supernatant fluid was decanted, the precipitate was washed with 0.5 ml 70% ethanol and dried in vacuo. After dissolving
55 the pellet in TE (10 mM Tris-HCl pH 7.4, 1 mM EDTA), the DNA content was measured by the ethidium bromide spot test, agarose plate method (Maniatis et al., Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory (1982) p. 468-469).

The ends of the linearized plasmid DNA were made blunt by incubating 30 ng for 2 hours at 15 °C in a 25 µl reaction mix containing 20 µM each of dATP, dGTP, dCTP, and TTP, 60 mM NaCl, 6 mM Tris-HCl pH 7.5, 6 mM MgCl₂, 1 mM dithiothreitol, and 22.5 units of DNA Polymerase I, large (Klenow) fragment (Boehringer-Mannheim). The reaction was terminated and the DNA purified by extraction with phenol/chloroform (Maniatis *et al.*, supra, p. 458-459) and ethanol precipitation (Maniatis *et al.*, supra, p. 461).

BamHI linkers (d-GGGATCCC, Boehringer-Mannheim), 12.5 µg, were phosphorylated with 40 units of T4 polynucleotide kinase (Pharmacia) in a 40 µl reaction mixture containing 50 mM Tris-HCl pH 7.4, 10 mM MgCl₂, 5 mM dithiothreitol, 500 µM ATP, and 40 µCi of γ -³²P-ATP (Amersham, 5000 Ci/mmol, 10 mCi/ml), for 30 minutes at 37 °C. The reaction was stopped by incubating at 70 °C for 5 minutes, and the linkers were stored at -20 °C until used.

The blunt-ended, linearized plasmid DNA was dissolved in 6.6 µl water and adjusted to a 10 µl final volume containing 125 ng phosphorylated BamHI linkers, 6.6 mM Tris-HCl pH 7.5, 6.6 mM MgCl₂, 1 mM ATP, 10 mM dithiothreitol, and 0.0025 units T4 DNA ligase (New England Biolabs). After incubating 18 hours at 4 °C, 5 µl of this mixture were added to 100 µl competent *E. coli* HB101 cells (BRL). Transformation of cells was performed according to the method provided by BRL. Eleven ampicillin-resistant colonies were chosen at random and each was used to inoculate 5 ml liquid culture of LB broth (Maniatis *et al.*, supra) containing 100 µg/ml ampicillin. After overnight growth at 37 °C, plasmid minipreps were made as described by Ish-Horowitz and Burke, *Nucleic Acids Research* 9:2989-2998 (1981).

By restriction enzyme mapping and agarose gel analysis, one plasmid, designated pJC220, was found to have a BamHI linker in place of the promoter-proximal HindIII site. This plasmid was also shown to retain the HindIII site (now unique) at the 3' end of the CheY coding region.

The pJC220 plasmid was converted to the pJC264 plasmid by digestion of 10 µg of pJC220 DNA with 50 units of HindIII (Boehringer-Mannheim) for 1 hour at 37 °C in a 50 µl solution containing 50 mM NaCl, 10 mM Tris-HCl pH 7.4, 10 mM MgSO₄, and 1 mM dithiothreitol. Ammonium acetate was added to 2.5 M final concentration, and the DNA recovered by precipitation with 2 volumes of ethanol. The HindIII digested DNA was then partially filled in with 5 units of the large fragment of DNA polymerase I (Boehringer-Mannheim) in a 20 µl solution containing 20 µM each dATP and dGTP, 60 mM NaCl, 6 mM Tris-HCl, pH 7.5, 6 mM MgCl₂, and 1 mM dithiothreitol, and incubated 30 minutes at room temperature. The sample was extracted with phenol/chloroform and recovered by ethanol precipitation as described by Maniatis *et al.*, *Molecular Cloning. A Laboratory Manual*. Cold Spring Harbor Laboratory (1982).

The DNA was dissolved in water and adjusted to 0.3 M NaCl, 30 mM Na acetate pH 4.6, and 4.5 mM ZnCl₂, in a final volume of 20 µl. Five units of SI nuclease (BRL) were added and the mixture incubated at 37 °C for 30 minutes. Digestion was stopped by adding 1 µl 0.5M EDTA pH 8.0, and the DNA was phenol/chloroform extracted and ethanol precipitated. The SI-nuclease treated DNA was digested with 80 units of EcoRI (New England Biolabs) in 50 µl of buffer containing 100 mM NaCl, 50 mM Tris-HCl pH 7.4, and 10 mM MgSO₄, for 30 minutes at 37 °C. DNA was recovered by ethanol precipitation in ammonium acetate as described above. The EcoRI ends were filled in with the large fragment of DNA polymerase I as described above, but on the presence of dATP and TTP and in the absence of dGTP and dCTP. DNA was extracted with phenol/chloroform and recovered by ethanol precipitation.

One hundred ng of this DNA were ligated for 24 hours at 4 °C in 10 µl of solution containing 66 mM Tris-HCl pH 7.5, 6.6 mM MgCl₂, 10 mM dithiothreitol, 1 mM ATP, and 400 units T4 DNA ligase (New England Biolabs). Two µl of ligation mix were used to transform 100 µl of competent *E. coli* JM109 cells (Stratagene) using the supplier's standard procedure. Ampicillin-resistant transformants were screened by colony hybridization using a 5'-³²P-labeled synthetic oligonucleotide d(CCCAAGAATTCAGTGG) as a probe, using standard methods of Mason & Williams, in "Nucleic Acid Hybridization: A Practical Approach, B.D. Hames and S.J. Higgins, eds. IRL Press (1985), p. 113-137. One hybridizing colony, designated pJC264, was shown by restriction mapping to have reconstructed a unique EcoRI site at the 3' end of the CheY gene.

The construction of pJC264 from CheY-ANF can be seen schematically in Figure 7 and the restriction map of pJC264 is shown in Figure 8.

EXAMPLE 12

Subcloning cDNA Inserts Into pJC264

Twenty micrograms of pJC264 from Example 11 was linearized with EcoRI using the reaction conditions described in Example 8. The reaction product was precipitated, washed twice with 70% EtOH

and suspended in 43 μ l of distilled water and 5 μ l of 10X CIP buffer (0.5 M Tris-HCl, pH 9.0, 10 mM $MgCl_2$, 1 mM $ZnCl_2$, 10 mM spermidine). The 5'-phosphate from the EcoRI ends was removed with calf intestinal alkaline phosphatase (Boehringer-Mannheim). One microliter of enzyme (19 U/ μ l) was added to initiate the reaction at 37 °C for 30 minutes and then a second microliter was added for an equivalent length of time.

5 The reaction was stopped by the addition of 42.5 μ l distilled water, 2.5 μ l 20% sodium dodecyl sulfate (SDS), 10 μ l 10X STE (100 mM Tris-HCl, pH 8.0/1 M NaCl/10 mM EDTA) and heated at 68 °C for 15 minutes. The reaction mixture was then extracted twice with phenol:chloroform/isoamyl alcohol (48:48:2), twice with chloroform/isoamyl alcohol (24:1), and the final aqueous phase was passed through a 1 cc column bed of Sephadex G-25 (medium) equilibrated in TE by centrifugation at 1000 x g for 5 minutes at room temperature (spin column). The DNA was then precipitated as described earlier, washed twice with 10 70% EtOH, suspended in 50 μ l of TE and quantitated by absorbance at 260nm.

Approximately 100 ng of EcoRI linearized and phosphatased pJC264 was mixed with an equimolar amount of gel purified *Eimeria tenella* cDNA insert in a 20 μ l reaction mixture which, in addition, consisted of 66 mM Tris-HCl, pH 7.6, 5 mM $MgCl_2$, 5 mM dithiothreitol, 1 mM ATP. The reaction was initiated by the addition of 1 μ l of T4 DNA ligase (New England Biolabs, 200-400 U/ μ l) and proceeded at 14 °C for 12-16 15 hours.

A predetermined volume (3 ml per transformation reaction) of 2 x YT bacterial media (16 g bacto-tryptone/10 g yeast extract/5 g NaCl per liter) was inoculated with a single colony of *E. coli* JM83 and grown with vigorous mixing at 37 °C until it reached an optical density at 600 nm of 0.6. Bacteria were 20 collected by centrifugation at 1000 x g at 4 °C for 5 minutes and gently suspended in one-half of the original culture volume with sterile 50 mM $CaCl_2$. The suspension was kept on ice for 20 minutes and the bacterial cells were collected by centrifugation as above. The pellet was then gently suspended in one-tenth volume of sterile 50mM $CaCl_2$. The bacterial suspension was then kept at 4 °C for 16-24 hours.

The 20 μ l ligation reaction mixture was diluted to 100 μ l by the addition of 80 μ l of sterile TE, and 5 μ l 25 and 95 μ l aliquots were dispensed to sterile polypropylene tubes. Approximately 200 μ l of competent bacteria were added to each of the tubes containing the ligation reactions (as well as the appropriate ligation and transformation controls) and these were placed on ice for 40 minutes. After this, the bacteria were "heat-shocked" by incubation at 42 °C for 90 seconds. Each transformation tube was then plated onto a 2 x YT agar plate which contained ampicillin at a concentration of 50 mg/l for the selection of bacteria 30 harboring plasmids and for plasmid maintenance. Plates were incubated in an inverted position overnight at 37 °C.

Bacterial clones harboring plasmids were identified by their ability to grow on plates in the presence of drug selection. Single colonies were used to inoculate 5 ml of 2 x YT/AMP (i.e., 2 x YT media containing ampicillin at 50 mg/L) and these cultures were grown overnight at 37 °C with vigorous shaking. Approximately 35 1.5 ml of the culture was poured off into an Eppendorf tube and collected by centrifugation in an Eppendorf centrifuge for at least 1 minute; the remainder of the culture was stored at 4 °C and served as a genetic stock. The media above the bacterial pellet was aspirated off and the pellet was suspended by vortexing in 100 μ l of a cold, freshly prepared solution of 50 mM glucose, 10 mM EDTA, 25 mM Tris-HCl (pH 8.0), 4 mg ml^{-1} lysozyme. This mixture was incubated at room temperature for 5 minutes. Then 200 μ l 40 of a cold, freshly prepared solution composed of 0.2 N NaOH and 1% SDS was added to each tube, mixed gently by inversion, and put on ice for 5 minutes. To this mixture was added 150 μ l of a cold, freshly prepared solution containing 6 ml of 5 M potassium acetate, 1.15 ml of glacial acetic acid, 2.85 ml distilled water. The contents were gently mixed on a vortex mixture and this mixture was stored on ice for 5 minutes. The cellular debris was collected by centrifugation in an Eppendorf centrifuge for 10 minutes at 45 4 °C and the supernatant was extracted one time with phenol:chloroform/isoamyl alcohol (25:24:1). Plasmid DNA and cellular RNA were precipitated from the final aqueous phase with the addition of two volumes of room temperature 100% ethanol. A pellet was collected by centrifugation for 5 minutes at room temperature, the pellet was washed one time with 70% ethanol and then dried briefly. The nucleic acid pellet was then suspended in 50 μ l of TE containing 20 μ g of DNase-free RNase per ml and incubated for 15-30 50 minutes at 37 °C to quantitatively eliminate cellular RNA. Aliquots of 10 μ l were then cut to completion with EcoRI (approximately 20 units) in a buffer composed of 50 mM NaCl, 100 mM Tris-HCl (pH 7.5), 5 mM $MgCl_2$ at 37 °C for 60 minutes. The restriction enzyme reaction products were fractionated by agarose gel electrophoresis to identify those plasmids which contained the appropriate inserts. Those recombinant plasmids which contained the predicted EcoRI insert were then cut with a second restriction enzyme 55 (usually Pst I) to verify (i) that only a single copy of the insert was contained within the plasmid, and (ii) to score for orientation of the insert DNA with respect to the bacterial promoter. This was accomplished by removing a second 10 μ l aliquot from the remaining 40 μ l of RNase-digested bacterial nucleic acid and cutting it in a buffer composed of 100 mM NaCl, 10 mM Tris-HCl (pH 7.5), and 10 mM $MgCl_2$ with

approximately 20 units of PstI for 60 minutes at 37°C. Again, the restriction enzyme digests were resolved by agarose gel electrophoresis.

EXAMPLE 13

Production of Eimeria-CheY Fusion Proteins

An overnight culture of selected recombinant bacteria was prepared by inoculating 5 mls of broth with a single colony bacteria. The culture medium was 2 x YT (16 g tryptone, 10 g yeast extract, 10 g NaCl/liter) containing ampicillin (50 µg/ml). The overnight culture was used to inoculate 500 mls of 2 x YT containing ampicillin. The culture was grown at 37°C with aeration until mid-log growth was reached ($A_{650} 0.5$) at which point IPTG was added to a final concentration of 100 µM. The culture was grown for a further 3-4 hours at 37°C, chilled on ice, and centrifuged for 15 minutes at 4°C. The cells were washed once with PBS, then the bacteria were collected by centrifugation and were stored frozen at -70°C until needed. When needed, the bacterial pellets were thawed and suspended in 10 ml of 30 mM Tris-HCl, pH 8.0, 50 mM EDTA and 1 mM phenylmethylsulphonylfluoride (Buffer A). The suspension was sonicated on ice twice, each time for three minutes, using a Branson cell disrupter Model 350 (duty cycle 30, output control 4). The sonicate was clarified by centrifugation at 27000 x g for 45 minutes at 4°C. The supernatant fluid constituted the first supernatant. The pellet of insoluble material was washed in 10 ml of buffer A containing 0.1% w/v Triton X100. The suspension was stirred in an ice-bath for 30 minutes prior to centrifugation at 27,000 x g for 45 minutes and at 4°C. The supernatant fluid is designated the second supernatant. The pellet (P₂) was then washed twice in Buffer A and the wash discarded. Pellet (P₂) was suspended in 1.0 ml of 6 M guanidine-HCl containing 100 mM dithiothreitol and the suspension was incubated for 2 hours at 50°C. The suspension was diluted to 10 mls with 7 M urea and was clarified by centrifugation at 27000 x g for 45 minutes at 4°C. The supernatant fluid constituted the third supernatant. Different fusion proteins exhibited different solubility properties, some were found predominantly in the first supernatant, some in the second, and some (most commonly) were found in the third.

The SO6-CheY antigen (recombinant A antigen) was found in the first, second, and third supernatant. Material for in vivo testing was prepared from the third supernatant by ion exchange chromatography. A Trisacryl M-DEAE (LKB) column (5 mls) equilibrated in 0.025 M Tris-HCl, pH 8.5, 8 M urea was prepared. From the third supernatant, a 2 ml sample, containing 12 mg protein, was dialyzed against 100 ml of the above buffer, and was then applied to the column. The column was washed with one column volume of column buffer, prior to step-wise elution with column buffer containing 0.05 M, 0.1 M, 0.15 M, 0.2 M, 0.25 M, 0.3 M, 0.35 M, or 0.4 M NaCl. Each elution was performed with two column volumes. Eluates were tested for the presence of recombinant protein by SDS-PAGE and Western blotting, using rabbit anti-Fraction V. The SO6/CheY protein was found to elute in the 0.15 M and 0.20 M NaCl fractions. Fractions were pooled, dialyzed against 50 mM NH₄CO₃, and were freeze dried. The yield of protein from a 500 ml culture is approximately 3 mg.

The SO7-CheY fusion protein (recombinant B antigen) was found in the third supernatant. Further purification was obtained by chromatography on hydroxyapatite. A column of hydroxyapatite (6 ml bed volume; BioRad Labs; HPT grade) and equilibrated in 7 M urea, and the third supernatant was applied to the column. After washing the column with one bed volume of 7 M urea, the flow-through and wash were combined, concentrated to 10 ml on Amicon diafiltration membrane YM10, dialyzed against 50 mM NH₄HCO₃, and were freeze dried (including any precipitate that formed). The yield from a 500 ml culture was approximately 35 mg protein.

The SP54-CheY fusion protein (recombinant C antigen) was also found in the third supernatant. Further purification was unnecessary for in vivo testing. The yield of protein in the third supernatant from a 500 ml culture was approximately 170 mg.

The SO311-CheY fusion protein (recombinant H antigen) was also found in the third supernatant. Further purification was obtained by chromatography on hydroxyapatite. The column was prepared as described above, and the third supernatant applied. The column was developed with two bed volumes of 7 M urea, then 2 bed volumes of 7 M urea containing 10 mM, 20 mM, 40 mM, 80 mM, 160 mM or 320 mM sodium phosphate buffer, pH 6.5. Column eluates were tested for the presence of recombinant protein by SDS-PAGE and Western blotting, using rabbit anti-fraction V, rabbit anti-sporozoite serum, or recombinant eluted antibodies. The SO311/CheY protein was found in the 40 mM, 80 mM and 160 mM eluates. These eluates were pooled, concentrated, dialyzed and freeze-dried exactly as above. The yield from a 500 ml culture was approximately 5 mg protein.

The SO216-CheY fusion protein (recombinant F antigen) was also found in the third supernatant fluid. No further purification was necessary for in vivo testing. The yield from a 500 ml culture was approximately 30 mg protein.

5 EXAMPLE 14

Characterization of Recombinant- 10 Derived E. tenella Immunogens

Representative E. tenella immunogen clones, from Example 9, were subjected to nucleotide sequence analysis utilizing one or two of three standard methodologies. Some sequence analyses were determined using the chemical degradation method of Maxam and Gilbert, Methods in Enzymology, 65 (part 1): 497-559 (1980). More commonly, the nucleotide sequence was determined by the dideoxy chain termination technique, using denatured plasmid templates (plasmid pUC18, containing assorted subsequences of the E. tenella cDNAs) as described by Hattori and Sakaki, Analyt. Biochem. 152: 232-238 (1986). The third approach to nucleotide sequence determination was accomplished by subcloning the cDNA insert, or portions of it, into bacteriophage mp18 and sequencing secreted single-stranded recombinant phage templates using the standard dideoxy chain-termination sequencing methodology of Messing, Methods in Enzymology 101: 20-78 (1983). In addition to AMV reverse transcriptase and the Klenow fragment of DNA polymerase I, a modified T7 DNA polymerase has been employed, see Tabor and Richardson, Proc. Nat. Acad. Sci. USA 84: 4767-4771 (1987).

The amino acid sequences were deduced from the determined nucleotide sequences by combining the following information. Each of the cDNAs, see Example 8, in the phage expression vector γ gt11 was identified by polyclonal antisera, see Example 2, when expressed as a fusion protein with β -galactosidase. The nature of the covalent attachment of this fusion protein is shown in the following table.

30 **TABLE 5**

EcoRI Cloning Site

EcoRI

<u>Beta-galactosidase</u>		<u>E. tenella</u>	
5'	GCG GAA TTC		3'
	Ala Glu Phe		

This junction (and reading frame, cloning site) at the EcoRI cleavage site, is regenerated in each subsequent cloning event involving the entire cDNA irrespective of the subcloning vector, pUC18, mp18 or pJC264. Consequently, the reading frame can be unequivocally identified and the nucleotide sequence translated once the orientation of the insert in these three vectors is established. The orientation of the cDNA insert in plasmid, puc 18 and pJC264, or phage, mp18, vectors is accomplished by restriction enzyme mapping, see Example 9. Once asymmetric restriction enzyme recognition sequences are identified within the cDNA insert, insert orientation and transcriptional orientation can be unequivocally assigned when the recognition sequences are similarly predicted by the nucleotide sequence.

Group A clone nucleotide sequences and the resulting Group A immunogen amino acid sequences are exemplified by the representative clone SO67. This clone is entirely contained within the SO6 clone, see

Example 9. Of the approximately 870 nucleotides in this clone the first 162 nucleotides starting at the 5' end have been sequenced. The transcriptional orientation and therefore the correct reading frame can be deduced unambiguously based upon the location in the nucleotide sequence of restriction enzyme recognition sequences which are predicted by restriction enzyme mapping of the CheY-SO67 recombinant plasmid. The nucleotide sequence and the resulting 53 N-terminal amino acid sequence is shown in the following table.

TABLE 6

**N-Terminal Nucleotide And Deduced Amino Acid
Sequence of Group A Immunogen S067**

	10	20	30	40	50
	*	*	*	*	*
10	T TTA TTC CTT CGA TGC CTG GCG GCG TTG TTC ATC ATG TTC ATC ACG AGG CGC CTT CTG				
20	Leu Phe Leu Arg Cys Leu Ala Ala Leu Phe Ile Met Phe Ile Thr Arg Arg Leu Leu				
	10				
25	60	70	80	90	100
	*	*	*	*	*
	CTG CTG CGA TTC ACC GTT CCT ACC GTG CTT TGC TGC TGC AGC AGC AGC ANG TGC TCG				
30	Leu Leu Arg Phe Thr Val Pro Thr Val Leu Cys Cys Cys Ser Ser Ser XXX Cys Ser				
	20	30			
35	120	130	140	150	160
	*	*	*	*	*
	TCG ANG NAG AGC GCC GGG GCA GCA GAA GCA GCA GCA GCA GCT CG				
40	Ser XXX XXX Ser Ala Gly Ala Ala Glu Ala Ala Ala Ala Ala Ala				
	40	50			

An additional 221 nucleotide sequence has been obtained from the 3' end of the clone, see table 7 below, but the reading frame has not been deduced.

TABLE 7

3' Nucleotide Sequence of Group A Immunogen S067

50	1	CGAGTGGCTG GTTGACACCG GCAGGGTCTT CGCCGGCGGC GTTGCTAGCA TAGCCGACGG
	61	CTGCCGGCTC TTCGGAGCAG CAGTGGAGGG CGAGGGCAAC GCTGGGAAGA ACTCGTCAAG
	121	ACCAACTACC AAATTGAAGT CCCCCAGGAA GACGGAACCT CCATTTCAGT GGATTGCGAC
55	181	GAGGCGGAGA CTCTGCGGCA GGCGGTGGTG GACGGCCGCG C

Group B clone nucleotide sequence and the resulting Group B immunogen amino acid sequence are exemplified by the representative clone S07. The reading frame can be deduced unambiguously by

correlating the position of restriction enzymes sites asymmetrically located within the cDNA with the location of their respective recognition sequences as predicted by the nucleotide sequence analysis. All 957 nucleotides in this clone have been sequenced. The nucleotide sequence and the amino acid sequence up to the termination codon at base 713 are shown in the following table.

TABLE 8
Nucleotide And Deduced Amino Acid Sequence
of Group B Immunogen S07

	10		20		30		40		50										
	*		*		*		*		*										
T	CTC	GCC	CCA	ACT	TTT	TCC	CCC	GCG	CTC	CGC	AGC	AGC	AGC	AGC	AGC	AGC	AGC	AGC	AGC
	Leu	Ala	Pro	Thr	Phe	Ser	Pro	Ala	Leu	Arg	Ser	Ser	Ser	Ser	Ser	Ser	Ser	Ser	Ser
	10																		
	60		70		80		90		100		110								
	*		*		*		*		*		*								
	AGC	AAA	ATG	GCA	GAC	CTC	TTC	AGC	GGA	CTC	GTG	GGC	GGC	GTC	GTC	GGC	GCT	GTT	GCT
	Ser	Lys	Met	Ala	Asp	Leu	Phe	Ser	Gly	Leu	Val	Gly	Gly	Val	Val	Gly	Ala	Val	Ala
	20																		
	30																		
	120		130		140		150		160		170								
	*		*		*		*		*		*								
	GCA	GCA	GAT	TTG	CCT	GCG	GAG	GGC	GAG	AGG	GCC	CCC	CGC	CCC	GCC	CCC	GGC	ACT	GCC
	Ala	Ala	Asp	Leu	Pro	Ala	Glu	Gly	Glu	Arg	Ala	Pro	Arg	Pro	Ala	Pro	Gly	Thr	Ala
	40																		
	50																		
	180		190		200		210		220										
	*		*		*		*		*										
	TGG	ACT	TGC	TGC	TGC	AGC	AAA	CTG	CAA	GAA	GGG	GCC	CGC	GAG	CTG	GAG	GGT	TTT	GTG
	Trp	Thr	Cys	Cys	Cys	Ser	Lys	Leu	Gln	Glu	Gly	Ala	Arg	Glu	Leu	Glu	Gly	Phe	Val
	60																		
	70																		
	230		240		250		260		270		280								
	*		*		*		*		*		*								
	CAG	CAG	CTG	AGT	TTT	GTT	GCA	GGG	AAG	CTG	GCC	TGC	TGC	CTG	CGG	GTG	GGG	GCG	GAG
	Gln	Gln	Leu	Ser	Phe	Val	Ala	Gly	Lys	Leu	Ala	Cys	Cys	Leu	Arg	Val	Gly	Ala	Glu
	80																		
	90																		
	290		300		310		320		330		340								
	*		*		*		*		*		*								
	CAG	CTG	GCG	CGC	TGC	GCT	GCG	GAG	GGG	CGG	CTG	CCC	AGC	AGC	AGC	AGC	AGC	AGC	AGC
	Gln	Leu	Ala	Arg	Cys	Ala	Ala	Glu	Gly	Arg	Leu	Pro	Ser	Ser	Ser	Ser	Ser	Ser	Ser
	100																		
	110																		
	350		360		370		380		390		400								
	*		*		*		*		*		*								
	TGC	TGC	GCG	CTG	CTG	CAG	CTC	GAG	AAG	CAG	GAC	CTC	GAG	CAG	AGC	CTC	GAG	GCC	GGC
	Cys	Cys	Ala	Leu	Leu	Gln	Leu	Glu	Lys	Gln	Asp	Leu	Glu	Gln	Ser	Leu	Glu	Ala	Gly
	120																		
	130																		

410 420 430 440 450
 * * * * *
 AAG CAG GGC GCG GAG TGC CTC TTG AGG AGC AGC AAA CTG GCC CTC GAG GCC CTC CTC
 Lys Gln Gly Ala Glu Cys Leu Leu Arg Ser Ser Lys Leu Ala Leu Glu Ala Leu Leu
 5 140 150

460 470 480 490 500 510
 * * * * * *
 GAG GGG GCC CGC GTT GCA GCA ACG CGG GGT TTG CTG CTG GTC GAG AGC AGC AAA GAC
 Glu Gly Ala Arg Val Ala Ala Thr Arg Gly Leu Leu Leu Val Glu Ser Ser Lys Asp
 10 160 170

520 530 540 550 560 570
 * * * * * *
 ACG GTG CTG CGC AGC ATT CCC CAC ACC CAG GAG AAG CTG GCC CAG GCC TAC AGT TCT
 Thr Val Leu Arg Ser Ile Pro His Thr Gln Glu Lys Leu Ala Gln Ala Tyr Ser Ser
 15 180 190

580 590 600 610 620
 * * * * *
 TTC CTG CGG GGC TAC CAG GGG GCA GCA GCG GGG AGG TCT CTG GGC TAC GGG GCC CCT
 Phe Leu Arg Gly Tyr Gln Gly Ala Ala Ala Gly Arg Ser Leu Gly Tyr Gly Ala Pro
 20 200

630 640 650 660 670 680
 * * * * * *
 GCT GCT GCT TAC GGC CAG CAG CAG CAG CCC AGC AGC TAC GGG GCG CCC CCC GCC TCC
 Ala Ala Ala Tyr Gly Gln Gln Gln Gln Pro Ser Ser Tyr Gly Ala Pro Pro Ala Ser
 25 210 220

690 700 710 720 730 740
 * * * * * *
 AGC CAG CAG CCC TCC GGC TTC TTC TGG TAG CCC TGC AGC AGC AGC AGC AGC AGC AGC
 Ser Gln Gln Pro Ser Gly Phe Phe Trp —
 30 230

750 760 770 780 790
 * * * * *
 AGC AGC AGC AGC GCG GGC GGC AGC CGC GGC GGG GCC GGG GCG CCG CTG CAG CAA CAG
 35 40

800 810 820 830 840 850
 * * * * * *
 CAG CAG CCG nnn CGG CTA GCG CCG CGG AGC ACT CGC AGG GAA CTC CAC AGG CAG CGG
 45

860 870 880 890 900 910
 * * * * * *
 GAG AGC AGC AGG GAC GAG AAG CAG GTC ATG TAG CGC AGG CAG CAG CGC CAG CTG CAG
 50

920 930 940 950
 * * * *
 CAG CAG CAG CAG CAG CAG CAG CAG CAG CAG CTC CTG CAC CG
 55

Group C clone nucleotide sequence and the resulting Group C immunogen amino acid sequence are exemplified by the representative clone SP54, see Example 9. This clone is entirely contained within the SP59 clone, see Example 9. Of the approximately 700 nucleotides in this clone the first 157 nucleotides

starting at the 5' end have been sequenced. The transcriptional orientation and therefore the appropriate reading frame can be unequivocally deduced by correlating restriction enzyme recognition sequences in the nucleotide sequence with their asymmetric location predicted by restriction enzyme mapping of the CheY-SP54 recombinant plasmid. The nucleotide sequence and the resulting 52 amino acid sequence is shown in the following table.

TABLE 9
N-Terminal Nucleotide and Deduced Amino Acid
Sequence of Group C Immunogen SP54

	10	20	30	40	50
	*	*	*	*	*
C	GCG GAA TCC GCA GAC ACT GCT GAG ATC CGC GTG CCC GTG GGG GCC ACT GTG GTG GTG				
	Ala Glu Ser Ala Asp Thr Ala Glu Ile Arg Val Pro Val Gly Ala Thr Val Val Val				
	10				
60	70	80	90	100	110
*	*	*	*	*	*
CGG CTT CAG AGC GTT GGG GGC TAC AGG CCA GTG TTG GTG AGT GCC CAG AGT GGG GCT					
Arg Leu Gln Ser Val Gly Gly Tyr Arg Pro Val Leu Val Ser Ala Gln Ser Gly Ala					
20	30				
120	130	140	150		
*	*	*	*		
GTG GGC CTC TCC GAG CTT TCC CAG GCT TCC CCC AGT TCG GCC					
Val Gly Leu Ser Glu Leu Ser Gln Ala Ser Pro Ser Ser Ala					
40	50				

Group H clone nucleotide sequence and the resulting Group H immunogen amino acid sequence is exemplified by the representative clone SO311, see Example 9. Of the approximately 650 nucleotides in this clone, the first 185 nucleotides starting at the 5' end have been sequenced. The transcriptional orientation and therefore the appropriate reading frame can be unequivocally deduced by correlating restriction enzyme recognition sequences in the nucleotide sequence with their asymmetric location predicted by restriction enzyme mapping. The nucleotide sequence and the resulting 61 amino acid sequence is shown in the following table.

TABLE 10

**N-Terminal Nucleotide and Deduced Amino Acid
Sequence of Group H Immunogen S0311**

5

10

15

20

25

30

35

40

10

20

30

40

50

60

70

80

90

100

110

120

130

140

150

160

170

180

190

200

210

220

230

240

250

260

270

280

290

300

310

320

330

340

350

360

370

380

390

400

410

420

430

440

450

460

470

480

490

500

510

520

530

540

550

560

570

580

590

600

610

620

630

640

650

660

670

680

690

700

710

720

730

740

750

760

770

780

790

800

810

820

830

840

850

860

870

880

890

900

910

920

930

940

950

960

970

980

990

1000

1010

1020

1030

1040

1050

1060

1070

1080

1090

1100

1110

1120

1130

1140

1150

1160

1170

1180

1190

1200

1210

1220

1230

1240

1250

1260

1270

1280

1290

1300

1310

1320

1330

1340

1350

1360

1370

1380

1390

1400

1410

1420

1430

1440

1450

1460

1470

1480

1490

1500

1510

1520

1530

1540

1550

1560

1570

1580

1590

1600

1610

1620

1630

1640

1650

1660

1670

1680

1690

1700

1710

1720

1730

1740

1750

1760

1770

1780

1790

1800

1810

1820

1830

1840

1850

1860

1870

1880

1890

1900

1910

1920

1930

1940

1950

1960

1970

1980

1990

2000

2010

2020

2030

2040

2050

2060

2070

2080

2090

2100

2110

2120

2130

2140

2150

2160

2170

2180

2190

2200

2210

2220

2230

2240

2250

2260

2270

2280

2290

2300

2310

2320

2330

2340

2350

2360

2370

2380

2390

2400

2410

2420

2430

2440

2450

2460

2470

2480

2490

2500

2510

2520

2530

2540

2550

2560

2570

2580

2590

2600

2610

2620

2630

2640

2650

2660

2670

2680

2690

2700

2710

2720

2730

2740

2750

2760

2770

2780

2790

2800

2810

2820

2830

2840

2850

2860

2870

2880

2890

2900

2910

2920

2930

2940

2950

2960

2970

2980

2990

3000

3010

3020

3030

3040

3050

3060

3070

3080

3090

3100

3110

3120

3130

3140

3150

3160

3170

3180

3190

3200

3210

3220

3230

3240

3250

3260

3270

3280

3290

3300

3310

3320

3330

3340

3350

3360

3370

3380

3390

3400

3410

3420

3430

3440

3450

3460

3470

3480

3490

3500

3510

3520

3530

3540

3550

3560

3570

3580

3590

3600

3610

3620

3630

3640

3650

3660

3670

3680

3690

3700

3710

3720

3730

3740

3750

3760

3770

3780

3790

3800

3810

3820

3830

3840

3850

3860

3870

3880

3890

3900

3910

3920

3930

3940

3950

3960

3970

3980

3990

4000

4010

4020

4030

4040

4050

4060

4070

4080

4090

4100

4110

4120

4130

4140

4150

4160

4170

4180

4190

4200

4210

4220

4230

4240

4250

4260

4270

4280

4290

4300

4310

4320

4330

4340

4350

4360

4370

4380

4390

4400

4410

4420

4430

4440

4450

4460

4470

4480

4490

4500

4510

4520

4530

4540

4550

4560

4570

4580

4590

4600

4610

4620

4630

4640

4650

4660

4670

4680

4690

4700

4710

4720

4730

4740

4750

4760

4770

4780

4790

4800

4810

4820

4830

4840

4850

4860

4870

4880

4890

4900

4910

4920

4930

4940

4950

4960

4970

4980

4990

5000

5010

5020

5030

5040

5050

5060

5070

5080

5090

5100

5110

5120

5130

5140

5150

5160

5170

5180

5190

5200

5210

5220

5230

5240

5250

5260

5270

5280

5290

5300

5310

5320

5330

5340

5350

5360

5370

5380

5390

5400

5410

5420

5430

5440

5450

5460

5470

5480

5490

5500

5510

5520

5530

5540

5550

5560

5570

5580

5590

5600

5610

5620

5630

5640

5650

5660

5670

5680

5690

5700

5710

5720

5730

5740

5750

5760

5770

5780

5790

5800

5810

5820

5830

5840

5850

5860

5870

5880

5890

5900

5910

5920

5930

5940

5950

5960

5970

5980

5990

6000

6010

6020

6030

6040

6050

6060

6070

6080

6090

6100

6110

6120

6130

6140

6150

6160

6170

6180

6190

6200

6210

6220

6230

6240

6250

6260

6270

6280

6290

6300

6310

6320

6330

6340

6350

6360

6370

6380

6390

6400

6410

6420

6430

6440

6450

6460

6470

6480

6490

6500

6510

6520

6530

6540

6550

6560

6570

6580

6590

6600

6610

6620

6630

6640

6650

6660

6670

6680

6690

6700

6710

6720

6730

6740

6750

6760

6770

6780

6790

6800

6810

6820

6830

6840

6850

6860

6870

6880

6890

6900

6910

6920

6930

6940

6950

6960

6970

6980

6990

7000

7010

7020

7030

7040

7050

7060

7070

7080

7090

7100

7110

7120

7130

7140

7150

7160

7170

7180

7190

7200

7210

7220

7230

7240

7250

7260

7270

7280

7290

7300

7310

7320

7330

7340

7350

7360

7370

7380

7390

7400

7410

7420

7430

7440

7450

7460

7470

7480

7490

7500

7510

7520

7530

7540

7550

7560

7570

7580

7590

7600

7610

7620

7630

7640

7650

7660

7670

7680

7690

7700

7710

7720

7730

7740

7750

7760

7770

7780

7790

7800

7810

7820

7830

7840

7850

7860

7870

7880

7890

7900

7910

7920

7930

7940

7950

7960

7970

7980

7990

8000

8010

8020

8030

8040

8050

8060

8070

8080

8090

8100

8110

8120

8130

8140

8150

8160

8170

8180

8190

8200

8210

8220

8230

8240

8250

8260

8270

8280

8290

8300

8310

8320

8330

8340

8350

8360

8370

8380

8390

8400

8410

8420

8430

8440

8450

8460

8470

8480

8490

8500

8510

8520

8530

8540

8550

8560

8570

8580

8590

8600

8610

8620

8630

8640

8650

8660

8670

8680

8690

8700

8710

8720

8730

8740

8750

8760

8770

8780

8790

8800

8810

8820

8830

8840

8850

8860

8870

8880

8890

8900

8910

8920

8930

8940

8950

8960

8970

8980

8990

9000

9010

9020

9030

9040

9050

9060

9070

9080

9090

9100

9110

9120

9130

9140

9150

9160

9170

9180

9190

9200

9210

9220

9230

9240

9250

9260

9270

9280

9290

9300

9310

9320

9330

9340

9350

9360

9370

9380

9390

9400

9410

9420

9430

9440

9450

9460

9470

9480

9490

9500

9510

9520

9530

9540

9550

9560

9570

9580

9590

9600

9610

9620

9630

9640

9650

9660

9670

9680

9690

9700

9710

9720

9730

9740

9750

9760

9770

9780

9790

9800

9810

9820

9830

9840

9850

9860

9870

9880

9890

9900

9910

9920

9930

9940

9950

9960

9970

9980

9990

10000

10010

10020

10030

10040

10050

10060

10070

10080

10090

10100

10110

10120

10130

10140

10150

10160

10170

10180

10190

10200

10210

10220

10230

10240

10250

10260

10270

10280

10290

10300

10310

10320

10330

10340

10350

10360

10370

10380

10390

10400

10410

10420

10430

10440

10450

10460

10470

10480

10490

10500

10510

10520

10530

10540

10550

10560

10570

10580

10590

10600

10610

10620

10630

10640

10650

10660

10670

10680

10690

10700

10710

10720

10730

10740

10750

10760

10770

10780

10790

10800

10810

10820

10830

10840

10850

10860

10870

10880

10890

10900

10910

10920

10930

10940

10950

10960

10970

10980

10990

11000

11010

11020

11030

11040

11050

11060

11070

11080

11090

11100

11110

11120

11130

11140

11150

11160

11170

11180

11190

11200

11210

11220

11230

11240

11250

11260

11270

11280

11290

11300

11310

11320

11330

11340

11350

11360

11370

11380

11390

11400

11410

11420

11430

11440

11450

11460

11470

11480

11490

11500

11510

11520

11530

11540

11550

11560

11570

11580

11590

11600

11610

11620

11630

11640

11650

11660

11670

11680

11690

11700

11710

11720

11730

11740

11750

11760

11770

11780

11790

11800

11810

11820

11830

11840

11850

11860

11870

11880

11890

11900

11910

11920

11930

11940

11950

11960

11970

11980

11990

12000

12010

12020

12030

12040

12050

12060

12070

12080

12090

12100

12110

12120

12130

12140

12150

12160

12170

12180

12190

12200

12210

12220

12230

12240

12250

12260

12270

12280

12290

12300

12310

12320

12330

12340

12350

12360

12370

12380

12390

12400

12410

12420

12430

12440

12450

12460

12470

12480

12490

12500

12510

12520

12530

12540

12550

12560

12570

12580

12590

12600

12610

12620

12630

12640

12650

12660

12670

12680

12690

12700

12710

12720

12730

12740

12750

12760

12770

12780

12790

12800

12810

12820

12830

12840

12850

12860

12870

12880

12890

12900

12910

12920

12930

12940

12950

12960

12970

12980

12990

13000

13010

13020

13030

13040

13050

13060

13070

13080

13090

13100

13110

13120

13130

13140

13150

13160

13170

13180

13190

13200

13210

13220

13230

13240

13250

13260

13270

13280

13290

13300

13310

13320

13330

13340

13350

13360

13370

13380

13390

13400

13410

13420

13430

1344

An additional 283 nucleotide sequence has been obtained from the 3' end of the clone, see table below, but the reading frame has not been deduced. Linker nucleotides are included in positions 1-8.

TABLE 11
3' Terminal Nucleotide Sequence of Group H
Immunogen SO311

1 GAATTCGGGT TATCCACATC ACGETGGACG TCTGATTTAG CGGAGGAGGT ATGAACCTC
 61 AGAGCCAGCC CAGTAGGAAG CATTATCCA TCTTGGTCTT TGCTCCACA GACGGTGACG
 121 GATTTCGAGG AGAGAGTGTA TCATTCTCT CAGTGTGGG ATGACATTCT CAGATGCGCG
 181 CATCACGTAA TGATAGCCAT TCCTGCTCCA GTCGGAAGCT ATGTCCTGAC TCTGGAGAGC
 241 AGCATTTCGG CGTGATACTT GAGCTTGTC GAGATAGCCA GCTGCTTCGA G

Group F clone nucleotide sequence of the Group F immunogen is exemplified by the representative clone SO216, see Example 9. The approximately 487 nucleotides, including eight linker nts at each end, have been sequenced. The sequence is given in the following table.

TABLE 12
Nucleotide Sequence of Group F Immunogen SO216

1 GAATTCGGGC AGAAAACAACT TACTGAAAGA CGGAGGGAAA GTGTCTCGCC GGCAAAGTTA
 61 AGCGAACGGA CTGATTTGGA AATAGGGTCT TGCTGCGCAA ACGAATGCTG CAAATGCATC
 121 CCAAAGCGGT ACCGCGATGG ATCAGCAAGA AAAACNCCTC AGTGAAACGA TAGGAGCTGA
 181 TGCCGAAGTC CGCACAGCAT GATCTATGTC TCATCGCTGC TGAGTTAGCT ACTGAGGCCA
 241 CACGGAAGGA GTGCTTTAGT TGTAGTTCTT GAGGTCTTCT ACGTGTACGG CATAGTCGAT
 301 GCTAGGGAAA CGAACAAGAG GGGCACCAGG TGACGACTCG TCGATGTACG CATGGAAGCC
 361 AGCAGCCGCC AGGACAGGCG TCAAGGCAAC GAGTGGGAGT AAAGCTTCAA TGGCGCTGTC
 421 TTTGCTGACT TTCGAGATCC AGGAGGTCTC GGCAGACTCG CTGACGGACT GGAGCAGCTC
 481 CGAATTC

The molecular weights of the primary *in vitro* translation products directed by mRNA specific for immunogens A, B, C and H were determined. *In vitro* translation of mRNA extracted from unsporulated oocysts, sporulating oocysts and sporozoites was performed using the rabbit reticulocyte cell free translation system, with either ³⁵S-methionine or ³H-leucine as the incorporated indicator isotope. Specific *in vitro* translation products were immuno-precipitated using monospecific antibodies, prepared as described in Example 6. The protocol for *in vitro* translation was as described in the technical bulletin from Promega Biotec (according to manufacturer's instructions) and for immunoprecipitation as in Taylor et al., Mol. Biochem. Parasitol. 10:305-318 (1983). The group A primary translation product recognized by monospecific antibody has a molecular weight of 24 kilo Daltons (kD). The major group B immunogen from clone S07 has a molecular weight of 28 kD while the minor immunogens have molecular weights of 170, 24, 22, 16, and 12kD. The additional minor specifically immunoprecipitable *in vitro* translation products were detectable when ³H-leucine was used as the labelled precursor amino acid. The 170 kD plus 22 kD minor immunogens were also detectable with ³⁵S-methionine. The major 28 kD immunogen was detectable only when ³H-leucine was used as the precursor amino acid. The molecular weight for the group C immunogen was not determined. The major group H immunogen from clone SO311 has a molecular weight of 28kD while the minor immunogens have molecular weights of 48, 38, 33, 16, 13, 12 and 10 kD. The additional minor specifically immunoprecipitable *in vitro* translation products were detectable when ³⁵S-methionine was used as the labelled precursor amino acid. The major 28 kD immunogen was detectable when both ³⁵S-methionine and ³H-leucine were used.

The specific mRNAs extracted from unsporulated and sporulating oocysts and/or sporozoites of *E. tenella*, Example 5, were sized by Northern blot analysis according to the method of Maniatis et al., Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, pg. 202 (1982) and the method described in Transfer and Immobilization of Nucleic Acids to S & S Solid supports, published by Schleicher and Schuell, Inc., pgs. 16-19 (1987). The mRNA complimentary to group A clone SO67, was 2.15 ± 0.13 kilobases (kb), to group B, clone S07, was 1.23 ± 0.22 kb; to group C, clones SP54 and SP59, was 1.12 ± 0.08 kb; and to group H, clone SO311, was 0.98 ± 0.07 kb.

Molecular weights and isoelectric points of *E. tenella* immunogens were also determined. Molecular weights were determined by analytical sodium dodecyl sulphate (SDS) polyacrylamide gel electrophoresis (PAGE) of samples prepared from sporulated oocysts and/or sporozoites of *E. tenella*, followed by transfer to nitrocellulose and immunodetection by Western Blotting as described in Example 6. Isoelectric points were determined by Western blotting of two dimensional gels of samples as described above. The dimensional gels were run according to the procedure of O'Farrell, J. Biol. Chem. 250: 4007-4021 (1975). Antibodies for both procedures were prepared as stated in Examples 2 and 6. The results are shown in the following table.

TABLE 13

Molecular Weight and Isoelectric Points of Native <i>E. tenella</i> Immunogens			
Immunogen group	Representative clones	Molecular weight (kD)	Isoelectric point
A	S06, S067	24	3.65
B	S07	27-28 22, 19, 18, 14, 12, 9, 6	5.1-6.3
C	SP54, SP59	21-22	n.d.
H	S0311	28, 18 27, 24, 23, 17 14, 12, 9	6.65
F	S0216	26-29	n.d.

The predominant B immunogen is characterized as a diffuse doublet of 27-28 kD on SDS-PAGE with the minor immunogens appearing as faint bands suggesting some sharing of antigenic determinants within *E. tenella*. The 27-28 doublet produces multiple spots on isoelectric focusing, in the range between pH 5.1 and 6.3. The pIs of the faint additional bands detected by Western blotting were not determined.

EXAMPLE 15

Induction Of Protection To Challenge With *E. tenella* By Recombinant-Derived *E. tenella* Immunogens

Broiler pullets were immunized three times via the intramuscular route on days 2, 9 and 16 days of age with samples containing 10 μ g of the specific recombinant fusion immunogen, from Example 13, in phosphate buffered saline absorbed on alum, 0.4% final concentration, in a total volume of 0.12 ml per dose per bird. The immunogen-alum complex was prepared by the procedure of Weir, Handbook of Experimental Immunology, Blackwell Scientific Publications London, pg. A3.11 (1978). Experimental and control birds were challenged at day 23, seven days after the final immunization, with an oral inoculation of from 5 to 30 x 10³ sporulated oocysts, an amount sufficient to yield a mean lesion score of at least 2.5 in non-immunized controls at 30 days of age. Seven days after challenge the chickens were killed and the severity of the lesions in the ceca was determined according to the method of Johnson and Reid, Exp. Parasitol. 28:30-36 (1970). Representative examples of the results are shown in Tables 14-18.

TABLE 14

Protection of Chickens Against Coccidiosis With Group A Immunogen SO67-CheY		
Challenge dose ($\times 10^{-3}$)	Immunized infected	Non-Immunized infected
5	2.18	3.41
10	2.57	3.57
15	1.78	3.44

TABLE 15

Protection of Chickens Against Coccidiosis With Group B Immunogen SO7-CheY		
Challenge dose ($\times 10^{-3}$)	Immunized infected	Non-Immunized infected
10	1.41	3.00
20	1.28	3.43
30	1.34	3.38

TABLE 16

Protection of Chickens Against Coccidiosis With Group C Immunogen SP54-CheY		
Challenge dose ($\times 10^{-3}$)	Immunized infected	Non-Immunized infected
5	1.71	3.38
10	1.68	3.00
15	1.93	3.22

TABLE 17

Protection of Chickens Against Coccidiosis With Group H Immunogen SO311-CheY		
Challenge dose ($\times 10^{-3}$)	Immunized infected	Non-Immunized infected
10	2.03	2.97
15	2.00	3.32

TABLE 18

Protection of Chickens Against Coccidiosis With Group F Immunogen SO216-CheY		
Challenge dose ($\times 10^{-3}$)	Immunized infected	Non-Immunized infected
10	1.50	2.16
15	1.30	2.72
20	1.25	2.89

These results show that recombinant *E. tenella* immunogens A, B, C, H, and F can be used to immunize two-day-old chickens against coccidiosis. Three intramuscular inoculations provide a high level of protection against the disease as indicated by the absence of severe lesion development in immune birds after a normally virulent infection.

EXAMPLE 16

Isolation Of The Native Form Of The B Immunogen From *Eimeria Tenella*

A suspension of 1×10^9 sporulated oocysts of *E. tenella* in 20 ml of phosphate buffered saline (PBS) containing 0.1 mM phenylmethylsulfonylfluoride (PMSF) was sonicated in an ice bath for a total of 10 minutes, in 2.5 minute bursts using a Branson Sonic Power Co. Sonifier Cell Disrupter 350 (duty cycle 30%, output control 4). The sonicate was centrifuged at $27,000 \times g$ for 30 minutes at 4°C . The pellet was washed 3 times in 40 ml PBS/0.1 mM PMSF, and was recovered by centrifugation as described above. The washed pellet was resuspended in 60 ml of 5 M guanidine-HCl/0.5 M Tris-HCl, pH 8.6, and 400 mg dithiothreitol. Reduction was allowed to proceed for 3 hours at 20°C with mild agitation. Insoluble debris was removed by centrifugation as described above. The supernatant fluid, containing reduced and solubilized B antigen was concentrated by ultrafiltration (Ultrafilter PM-10, Amicon Corp.) to 20 ml, and iodoacetic acid (400 mg) was added. The pH was readjusted to 8.6 by the addition of 3 M Tris base, and carboxymethylation was allowed to proceed for 60 minutes at 20°C in the dark. The reaction mixture was then dialyzed for 48 hours against 0.05 M NH_4HCO_3 /0.1 mM PMSF/0.02% sodium azide. With the removal of guanidine-HCl, some insoluble material formed which was subsequently removed by centrifugation as described above. The cleaned supernatant was then concentrated to 12 ml by ultrafiltration, as described above. The concentrate was then applied to a sizing column of Sephacryl S-200 (87 x 2.5 cm) equilibrated in 0.05 M NH_4HCO_3 , 0.1% Zwittergent 3-12 (Calbiochem), 0.02% sodium azide. A total of 120 x 4.5 ml fractions were collected, at a flow rate of 25 ml/hour. Effluent fractions were monitored at 280 nm, and the elution of the B immunogen was monitored by Western blotting, initially using rabbit anti-sporozoite antiserum, and subsequently with a rabbit antiserum to the SO7/CheY protein. Fractions containing the B antigen (47-57) were pooled, concentrated to 10 ml, and were reappplied to the column. The column was eluted and monitored as before. Pooled fractions were concentrated to a volume containing approximately 0.5 mg protein/ml. The total yield was 5.8 mg.

SDS gel analysis showed a single homogenously pure protein of $30 \text{ kD} \pm 3 \text{ kD}$, which on Western blot analysis was reactive with both rabbit anti-sporozoite antiserum and rabbit anti-SO7-CheY.

The immunogenic activity of this sample of B antigen purified by *E. tenella* was measured as described in Example 15. Two day old broiler pullets were immunized three times via the intramuscular route on days 2, 9 and 16 with samples containing 10 μg of the purified native B immunogen absorbed on alum (0.4% final concentration). The immunogen-alum complex was prepared by the procedure of Weir, Handbook of Experimental Immunology, Blackwell Scientific Publications, London, pg. A3-11 (1978). Experimental and control birds were challenged at day 23, seven days after the final immunization with an oral inoculation of from $5\text{-}15 \times 10^3$ sporulated oocysts. Seven days after challenge, the chickens were killed and the severity of the lesions in the ceca was determined according to the method of Johnson and Reid, Exp. Parasitol, 28, 30-36 (1970). Results are presented as mean cecal lesion scores for groups of eight birds and are shown in the following table.

TABLE 19

Protection of Chickens Against Coccidiosis With Native Group B Immunogen		
Challenge Dose ($\times 10^{-3}$)	Immunized Infected	Non-Immunized Infected
5	1.36	3.41
10	1.64	3.57
15	1.54	3.44

An alternative method of purifying the B immunogen from *E. tenella* is by affinity chromatography, using the antibody to the SO7-CheY protein. For this purpose, two affinity columns were prepared, one using serum from a rabbit removed prior to immunization with the SO7-CheY antigen (prebleed column), and one using antiserum from the same rabbit immunized with the SO7-CheY antigen, using the immunization regime described in Example 2. The SO7-CheY immunogen was prepared as described in Example 13. The immunoglobulin IgG fraction was prepared from 4 ml of each serum, using the method of Corthier et al., J. Immunol. Met., 66, 75-79 (1984). For each column, 15 mg of IgG was coupled to 0.5 gm of Sepharose-Protein A (Sigma), using the method of Schneidert et al., J. Biol. Chem. 257, 10766-10769 (1982). Coupling efficiency was between 75-95%. For immunoaffinity purification, approximately 5 mg of the reduced,

carboxymethylated extract, prepared as described above (with no purification by gel filtration), on 0.1 M borate buffer, pH 8.1, 0.5 M NaCl, 0.02% NaN₃, 0.1 mM PMSF, was applied to the prebleed column equilibrated in the same buffer. The column was washed with 3 ml of column buffer, and the combined column flow-through and washes were then applied to the anti-SO7/CheY column equilibrated in the same buffer. The column was washed with 10 ml of column buffer, prior to elution with 3 M NaSCN. The eluate was dialyzed for 48 hours versus 0.05 M NH₄HCO₃, prior to freezing. A total of approximately 50 µg protein was recovered in the final eluate.

The immunogenic activity of this affinity purified B antigen from *E. tenella* was tested as described in Example 15. Two day old broiler pullets were immunized three times via the intramuscular route on days 2, 9 and 16 with samples containing approximately 0.3 µg of the immunoaffinity purified Group B immunogen absorbed on alum (0.4% final concentration). The immunogen-alum complex was prepared by the procedure of Weir, Handbook of Experimental Immunology, Blackwell Scientific Publications, London, pg. A3-11 (1978). Experimental and control birds were challenged at day 23, seven days after the final immunization, with an oral inoculation of from 10-30 x 10³ sporulated oocysts. Seven days after challenge, the chickens were killed, and the severity of the lesions in the ceca was determined according to the method of Johnson and Reid, Exp. Parasitol. 28, 30-36. Results are presented as mean cecal lesion scores for groups of eight birds and are shown in the following table.

TABLE 20

Protection of Chickens Against Coccidiosis With Native Group B Immunogen		
Challenge Dose (x10 ⁻³) oocysts	Immunized	Infected
10	1.41	3.00
20	1.44	3.43
30	1.59	3.38

EXAMPLE 17

IDENTIFICATION AND ISOLATION OF B ANTIGENS FROM OTHER EIMERIA SPECIES

Fimeria *Acervulina* antigens were prepared by resuspending sporulated oocysts at a concentration of 5.5 x 10⁷ per ml and DEAE-52 purified sporozoites at a concentration of 2.6 x 10⁸ per ml in NET buffer (10 mM Tris-HCl, pH 8.0, 150 mM NaCl, 5 mM EDTA) containing a cocktail of protease inhibitors (2 mg/ml 1-10 phenanthroline, 2 mg/ml benzamidine, 0.002 mg/ml PMSF, 0.048 mg/ml Sigma soybean trypsin inhibitor, 0.048 mg/ml aprotinin, 0.02 mg/ml leupeptin). At this point, the samples were mixed with an equal volume of 2X sample buffer (0.125 M Tris-HCl, pH 6.8, 4% v/v SDS, 10% v/v 2-mercaptoethanol, 20% glycerol). The samples were boiled for 3 minutes, sonicated till fully disrupted, and reboiled again for 3 minutes. Bromophenol blue was added to 0.0025% and the samples were stored at -20 °C until ready for use.

For immunoblotting, antigens obtained from 3 x 10⁵ sporulated oocysts and 2 x 10⁶ sporozoites were loaded per slot and subjected to electrophoresis on a 5-20% SDS-polyacrylamide gradient gel (Laemmli, Nature 227: 680-684, 1970). Proteins separated by SDS-PAGE were electrophoretically transferred to nitrocellulose by the technique of Towbin et al, Proc. Natl. Acad. Sci. USA 76: 4350-4354 (1979). The nitrocellulose was blocked with 0.5% gelatin in phosphate buffered saline, pH 7.4, for 1 hour with three 200 ml washes followed by a second blocking with 0.25% gelatin in TEN buffer (50 mM Tris-HCl, 150 mM NaCl, 5 mM EDTA, pH 7.4) for 1 hour and washed as before. After blocking, the nitrocellulose was incubated overnight at room temperature in 20 ml of antibody (raised against the CheY fusion protein containing the *E. tenella* clone SO7 sequence, representing the B antigen of *E. tenella*) diluted 1:100 with TEN buffer containing 0.25% gelatin and 0.05% Triton X-100. The filters were washed 5 times for 20 minutes each with 200 ml of TEN containing 0.25% gelatin. Bound antibody was detected with ¹²⁵I-protein A diluted in 20 ml of TEN, 0.25% gelatin, 0.05% Triton to a final concentration of 2 x 10⁵ cpm/ml. Incubation with radiolabeled protein A was carried out for 1 hour at room temperature, after which time the filters were washed 2 times for 15 minutes with 200 ml of TEN containing 0.25% gelatin and 0.05% Triton and 4 times for 15 minutes with 200 ml of TEN. After washing, the filters were air dried and exposed to Kodak X-omat AR film. The antigen had a molecular weight of approximately 26 kD ± 3 kD.

Eimeria Maxima sporulated oocysts stored in phosphate-buffered saline (PBS) pH 7.6 were centrifuged for 10 minutes at 1600 x g and were resuspended in a volume of PBS equal to the packed cell pellet. To this suspension, an equal volume of glass beads was added and the oocysts were broken by orbital shaking at 200 rpm., Dulski, P. and Turner, M., Avian Diseases 32: 235-239, 1988. The disrupted oocysts and sporocysts were collected by centrifugation at 1600 x g for 10 minutes. The pellet was then resuspended in 50% Percoll/IX PBS and recentrifuged. The pellet containing clean sporocysts was collected and washed 2 times with PBS before proceeding. Undisrupted oocysts were collected from the top of the Percoll gradient, washed, and the entire procedure was repeated when necessary. Approximately 1.5 to 2 sporocysts were obtained per sporulated oocyst.

Antigens were prepared by resuspending the sporocysts at a concentration of 5×10^7 per ml in NET buffer (10 mM Tris-HCl, pH 8.0, 150 mM NaCl, 5 mM EDTA) containing a cocktail of protease inhibitors (2 mg/ml 1-10 phenanthroline, 2 mg/ml benzamidine, 0.002 mg/ml PMSF, 0.048 mg/ml Sigma soybean trypsin inhibitor, 0.048 mg/ml aprotinin, 0.02 mg/ml leupeptin). At this point, the samples were mixed with an equal volume of 2X sample buffer (0.125 M Tris-HCl, pH 6.8, 4% v/v SDS, 10% v/v 2-mercaptoethanol, 20% glycerol). The samples were boiled for 3 minutes, sonicated till fully disrupted, and reboiled again for 3 minutes. Bromophenol blue was added to 0.0025% and the samples were stored at -20 °C until ready for use.

For immunoblotting, antigens obtained from 1×10^6 sporocysts were loaded per slot and subjected to electrophoresis on a 5-20% SDS-polyacrylamide gradient gel (Laemmli, Nature 227: 680-684, 1970). Proteins separated by SDS-PAGE were electrophoretically transferred to nitrocellulose by the technique of Towbin et al, Proc. Natl. Acad. Sci. USA 76: 4350-4354 (1979). The nitrocellulose was blocked with 0.5% gelatin in phosphate buffered saline, pH 7.4, for 1 hour with three 200 ml washes followed by a second blocking with 0.25% gelatin in TEN buffer (50 mM Tris-HCl, 150 mM NaCl, 5 mM EDTA, pH 7.4) for 1 hour and washed as before. After blocking, the nitrocellulose was incubated overnight at room temperature in 20 ml of antibody (raised against the CheY fusion protein containing the *E. tenella* clone SO7 sequence, representing the B antigen of *E. tenella*) diluted 1:100 with TEN buffer containing 0.25% gelatin and 0.05% Triton X-100. The filters were washed 5 times for 20 minutes each with 200 ml of TEN containing 0.25% gelatin. Bound antibody was detected with 125 I-protein A diluted in 20 ml of TEN, 0.25% gelatin, 0.05% Triton to a final concentration of 2×10^5 cpm/ml. Incubation with radiolabeled protein A was carried out for 1 hour at room temperature, after which time the filters were washed 2 times for 15 minutes with 200 ml of TEN containing 0.25% gelatin and 0.05% Triton and 4 times for 15 minutes with 200 ml of TEN. After washing, the filters were air dried and exposed to Kodak X-omat AR film. The antigen had a molecular weight of approximately $28 \text{ kD} \pm 3 \text{ kD}$.

Sporulated oocysts were furnished as a 20 ml suspension in PBS, containing approximately 7×10^9 oocysts/ml. The suspension was made with 0.1 mM in PMSF, and then sonicated in an ice-bath until by microscopic examination no more than about 10% of intact sporozoites remained. The sporulated oocyst sonicate insoluble fraction, which contains the B-antigen, was obtained by collecting the pellet following centrifugation at 30,000 x g for 45 minutes at 4 °C. The B-antigen was extracted from the sonicate pellet by reduction and carboxymethylation. Briefly, the pellet was washed three times in PBS, then suspended in 60 ml 5M guanidine-HCl/0.5M Tris-HCl, pH 8.6 at room temperature. The suspension was then charged with 400 mg dithiothreitol (DTT) and kept at room temperature for 24 hours with mild agitation. The suspension was then centrifuged at 30,000 x g for 60 minutes at 4 °C, and the supernatant fluid was concentrated to 20 ml by ultrafiltration on an Amicon YM 10 membrane. The concentrate was charged with a fourfold molar excess, with respect to DTT, of 630 mg iodoacetic acid. The pH was adjusted immediately to 8.6 with 3M Tris base, and the system was kept at room temperature and in the dark for 2 hours, followed by dialysis against 50 mM NH_4HCO_3 /0.1 mM PMSF/0.02% sodium azide for 48 hours, with several changes of the permeate. During dialysis, some insoluble matter formed which was removed by centrifugation at 30,000 x g for 30 minutes at 4 °C. The retentate constitutes the reduced and carboxymethylated soluble fraction (RCSF), containing on the average 400 mg protein/ml in a total of about 30 ml.

RCSF derived from *E. acervulina* was applied to a column (100 x 2 cm) of Sephacryl S-200 equilibrated in running buffer (50 mM NH_4HCO_3 /0.1% zwittergent/0.1 mM PMSF, 0.02% sodium azide). A total of 60 fractions (5.5 ml) were collected. Fractions containing the antigen were identified by SDS-PAGE followed by Western blotting, using antiserum raised against the CheY fusion protein containing the *E. tenella* clone SO7 sequence, representing the B antigen of *E. tenella*. Appropriate fractions were pooled and concentrated to ca. 12 ml by ultrafiltration (Amicon, YM10 membrane). The concentrate was then passed over an affinity matrix constructed with pre-bleed IgG isolated from serum of the rabbit subsequentially injected with the SO7 fusion protein. Passage was done by continuous recycling for 18 hours at 4 °C. The flow-through from this step, containing the B-antigen, was diafiltered (Amicon, YM10 membrane) with 50 mM sodium

bicarbonate buffer to remove the zwittergent, then with column wash (0.1 M sodium borate, pH 8.0/0.5 M NaCl, 0.02% sodium azide/0.1 mM PFST). Final volume was about 12 ml and this constituted the charge (14361-216-2) to the biospecific matrix containing the rabbit anti-SO7 IgG. The charge was applied to the matrix by recycling for 18 hours at 4°C.

After draining the column of the charge fluid, the matrix was washed with the following and in the order indicated:

2 x 5 ml column wash

2 x 5 ml 10 mM Tris-HCl, pH 8.0

The antigen was then desorbed from the matrix by gravity passage of 10 ml of 0.1% zwittergent 3-12 in 10 mM Tris-HCl, pH 8.0. The eluate was dialysed against 50 mM NH_4HCO_3 and concentrated to 5.0 ml by ultrafiltration (Amicon, YM10 membrane). The protein was characterized by SDS-PAGE and silver staining.

The purification of the B-antigen from *E. maxima* followed a similar protocol as described, but pre-purification on Sephacryl S-200 was omitted. RSCF from *E. maxima*, 20 ml, was sub-aliquoted (10 ml each). The aliquot was charged with noctyl glucoside to 0.2% and with NaCl to 0.5 M and was then passed by recycling over a pre-bleed column for 18 hours at 4°C. The flow-through from this step constituted the column charge to the biospecific matrix, constructed with anti-SO7 IgG. Washing of the matrix and desorption of the antigen therefrom, was done as described above. The eluate was diafiltered with 10 mM Tris-HCl, pH 8.0 (Amicon YM10 membrane) to remove zwittergent, and was reconstituted in 2.5 ml 4 M urea (final product 38893-49-3). The product was characterized by SDS-PAGE and silver staining.

Biospecific matrices were prepared based on the method of Bethel et al., see above. This 1,1-carbonyldiimidazole activated support is available commercially under the trade name Reactigel (Pierce Chem. co., Rockford, ILL.) the feature of this support rests in the formation of a very stable and uncharged N-alkyl carbamate bond on reaction with a free amino group on the ligand. Reactigel was used in accordance with the manufacturer's suggested procedure. Briefly, 5 cc of an acetone suspension (50% bed volume) of Reactigel was transferred into an Econocolumn (BioRad) and drained free of acetone. Immunoglobulin IgG was then introduced as a solution of 12 mg/8 ml coupling buffer (0.1 M NaHCO_3 /0.5 M NaCl, pH 8.5). The column was sealed, mounted on a rocking platform and kept at 4°C overnight. The column was drained, and washed with 5 ml coupling buffer. Quenching was done by suspending the support in 10 ml coupling buffer containing 50 ml aminoethanol and placing the column on a rocking platform for 4 hours at room temperature. The matrix was then washed with 10 ml coupling buffer, 6 ml 3.5 M sodium thiocyanate, and finally with 10 ml "column wash" (0.1 M borate buffer, pH 8.0, 0.5 M NaCl, 0.02% sodium azide, 0.1 mM PMSF).

Chickens were immunized with the Group B immunogen isolated from *E. acervulina* and challenged with sporulated oocysts. The results are shown in the following table.

TABLE 21

Protection of Chickens Against Coccidiosis With Native Group B Immunogen From <i>Eimeria Acervulina</i>				
Antigen	Mean Lesion Score Challenge Organism			
	<i>E. acervulina</i> *		<i>E. tenella</i> *	
	Trial 1	Trial 2	Trial 1	Trial 2
None	2.90	2.40	2.75	2.46
<i>E. tenella</i> Recombinant Group B	1.64	1.65	2.13	1.60
<i>E. acervulina</i> Group B 0.1 µg	2.17	1.90	1.61	1.32
<i>E. acervulina</i> Group B 0.3 µg	1.90	1.15	2.03	1.36

* Challenge doses: *E. acervulina* 1-2 x 10⁵, *E. tenella* 2-5 x 10⁴

Samples of the expression vector pJC264 containing the DNA for the various *E. tenella* immunogens have been deposited in a host *Escherichia coli*, JM83 or JM109, under the Budapest Treaty in the American Type Culture Collection, 12301 Parklawn Drive, Rockville, Maryland 20852, USA.

On November 4, 1987, a sample of the expression vector containing clone SO7 was deposited and was given accession number 67577.

Claims

- Claims for the following Contracting States : AT, BE, CH, LI, DE, FR, GB, IT, LU, NL

1. A recombinant DNA molecule comprising a nucleotide sequence corresponding to all or a portion of the following nucleotide sequence:

T CTC GCC CCA ACT TTT TCC CCC GCG CTC CGC AGC AGC AGC AGC AGC AGC AGC AGC AGC
 AGC AAA ATG GCA GAC CTC TTC AGC GGA CTC GTG GGC GGC GTC GTC GGC GCT GTT GCT
 10 GCA GCA GAT TTG CCT GCG GAG GGC GAG AGG GCC CCC CGC CCC GCC CCC GGC ACT GCC
 TGG ACT TGC TGC TGC AGC AAA CTG CAA GAA GGC GCC CGC GAG CTG GAG GGT TTT GTG
 CAG CAG CTG AGT TTT GTT GCA GGG AAG CTG GCC TGC TGC CTG CGG GTG GGG GCG GAG
 15 CAG CTG GCG CGC TGC GCT GCG GAG GGG CGG CTG CCC AGC AGC AGC AGC AGC AGC AGC
 TGC TGC GCG CTG CTG CAG CTC GAG AAG CAG GAC CTC GAG CAG AGC CTC GAG GCC GGC
 AAG CAG GGC GCG GAG TGC CTC TTG AGG AGC AGC AAA CTG GCC CTC GAG GCC CTC CTC
 GAG GGG GCC CGC GTT GCA GCA ACG CGG GGT TTG CTG CTG GTC GAG AGC AGC AAA GAC
 20 ACG GTG CTG CGC AGC ATT CCC CAC ACC CAG GAG AAG CTG GCC CAG GCC TAC AGT TCT
 TTC CTG CGG GGC TAC CAG GGG GCA GCA GCG GGG AGG TCT CTG GGC TAC GGG GCC CCT
 GCT GCT GCT TAC GGC CAG CAG CAG CAG CCC AGC AGC TAC GGG GCG CCC CCC GCC TCC
 25 AGC CAG CAG CCC TCC GGC TTC TTC TGG TAG CCC TGC AGC AGC AGC AGC AGC AGC AGC
 AGC AGC AGC AGC GCG GGC GGC AGC CGC GGC GGG GCC GGG GCG CCG CTG CAG CAA CAG
 CAG CAG CCG nnn CGG CTA GCG CCG CGG AGC ACT CGC AGG GAA CTC CAC AGG CAG CGG
 GAG AGC AGC AGG GAC GAG AAG CAG GTC ATG TAG CGC AGG CAG CAG CGC CAG CTG CAG
 30 CAG CAG CAG CAG CAG CAG CAG CAG CAG CAG CTC CTG CAC CG

which encodes a Group B immunogen of Eimeria tenella.

2. The recombinant DNA molecule of Claim 1 wherein the DNA sequence is directly preceeded by and operably attached to a second DNA nucleotide sequence resulting in a fused DNA molecule.
 3. The second DNA sequence of Claim 2 wherein the DNA sequence expresses a CheY protein and the product of linking nucleotides.
 4. A recombinant DNA molecule comprising a nucleotide sequence capable of being expressed as at least one protein displaying the immunogenicity of a Group B immunogen of Eimeria tenella which sequence is:

1 T CTC GCC CCA ACT TTT TCC CCC GCG CTC CGC AGC AGC AGC AGC AGC AGC AGC AGC AGC
 AGC AAA ATG GCA GAC CTC TTC AGC GGA CTC GTG GGC GGC GTC GTC GGC GCT GTT GCT
 5 GCA GCA GAT TTG CCT GCG GAG GGC GAG AGG GCC CCC CGC CCC GCC CCC GGC ACT GCC
 TGG ACT TGC TGC TGC AGC AAA CTG CAA GAA GGG GCC CGC GAG CTG GAG GGT TTT GIG
 CAG CAG CTG AGT TTT GTT GCA GGG AAG CTG GCC TGC TGC CTG CGG GTG GGG GCG GAG
 CAG CTG GCG CGC TGC GCT GCG GAG GGG CGG CTG CCC AGC AGC AGC AGC AGC AGC AGC
 10 TGC TGC GCG CTG CTG CAG CTC GAG AAG CAG GAC CTC GAG CAG AGC CTC GAG GCC GGC
 AAG CAG GGC GCG GAG TGC CTC TTG AGG AGC AGC AAA CTG GCC CTC GAG GCC CTC CTC
 GAG GGG GCC CGC GTT GCA GCA ACG CGG GGT TTG CTG CTG GTC GAG AGC AGC AAA GAC
 15 ACG GTG CTG CGC AGC ATT CCC CAC ACC CAG GAG AAG CTG GCC CAG GCC TAC AGT TCT
 TTC CTG CGG GGC TAC CAG GGG GCA GCA GCG GGG AGG TCT CTG GGC TAC GGG GCC CCT
 GCT GCT GCT TAC GGC CAG CAG CAG CAG CCC AGC AGC TAC GGG GCG CCC CCC GCC TCC
 AGC CAG CAG CCC TCC GGC TTC TTC TGG TAG CCC TGC AGC AGC AGC AGC AGC AGC AGC
 20 AGC AGC AGC AGC GCG GGC GGC AGC CGC GGC GGG GCC GGG GCG CCG CTG CAG CAA CAG
 CAG CAG CCG nnn CGG CTA GCG CCG CGG AGC ACT CGC AGG GAA CTC CAC AGG CAG CGG
 GAG AGC AGC AGG GAC GAG AAG CAG GTC ATG TAG CGC AGG CAG CAG CGC CAG CTG CAG
 25 CAG CAG CAG CAG CAG CAG CAG CAG CAG CAG CAG CTC CTG CAC CG.

5. The recombinant DNA molecule of Claim 4 wherein the DNA sequence is directly preceeded by and operably attached to a second DNA nucleotide sequence resulting in a fused DNA molecule.
6. The second DNA sequence of Claim 5 wherein the DNA sequence expresses a CheY protein and the product of linking nucleotides.
7. The recombinant DNA molecule as claimed in any one of Claims 1, 2 or 5 wherein the DNA sequence is contained in an expression vector capable of expressing all or a portion of the immunogen DNA when incorporated in a host cell.
8. A recombinant Eimeria tenella protein immunogen comprising an amino acid sequence expressed by the group B DNA molecule of claim 1 and any microheterogeneous or subunit immunogen forms thereof.

9. A Group B Eimeria tenella protein immunogen comprising at least the following amino acid sequence:

10

Leu-Ala-Pro-Thr-Phe-Ser-Pro-Ala-Leu-Arg-Ser-Ser-Ser-Ser-Ser-Ser-Ser-

20

30

Ser-Ser-Lys-Met-Ala-Asp-Leu-Phe-Ser-Gly-Leu-Val-Gly-Gly-Val-Val-Gly-Ala-

40

50

Val-Ala-Ala-Ala-Asp-Leu-Pro-Ala-Glu-Gly-Glu-Arg-Ala-Pro-Arg-Pro-Ala-Pro-

60

70

Gly-Thr-Ala-Trp-Thr-Cys-Cys-Cys-Ser-Lys-Leu-Gln-Glu-Gly-Ala-Arg-Glu-

80

Leu-Glu-Gly-Phe-Val-Gln-Gln-Leu-Ser-Phe-Val-Ala-Gly-Lys-Leu-Ala-Cys-Cys-

90

100

Leu-Arg-Val-Gly-Ala-Glu-Gln-Leu-Ala-Arg-Cys-Ala-Ala-Glu-Gly-Arg-Leu-Pro-

110

120

Ser-Ser-Ser-Ser-Ser-Ser-Ser-Ser-Cys-Cys-Ala-Leu-Leu-Gln-Leu-Glu-Lys-Gln-Asp-

130

140

Leu-Glu-Gln-Ser-Leu-Glu-Ala-Gly-Lys-Gln-Gly-Ala-Glu-Cys-Leu-Leu-Arg-Ser-

5

150

160

Ser-Lys-Leu-Ala-Leu-Glu-Ala-Leu-Leu-Glu-Gly-Ala-Arg-Val-Ala-Ala-Thr-Agr-

10

170

Gly-Leu-Leu-Leu-Val-Glu-Ser-Ser-Lys-Asp-Thr-Val-Leu-Arg-Ser-Ile-Pro-His-

180

190

15

Thr-Gln-Glu-Lys-Leu-Ala-Gln-Ala-Tyr-Ser-Ser-Phe-Leu-Arg-Gly-Tyr-Gln-Gly-

200

210

20

Ala-Ala-Ala-Gly-Arg-Ser-Leu-Gly-Tyr-Gly-Ala-Pro-Ala-Ala-Ala-Tyr-Gly-Gln-

220

230

25

Gln-Gln-Gln-Pro-Ser-Ser-Tyr-Gly-Ala-Pro-Pro-Ala-Ser-Ser-Gln-Gln-Pro-Ser-

Gly-Phe-Phe-Trp

30

and microheterogeneous or subunit immunogen forms thereof.

10. The Group B Eimeria tenella immunogen of Claim 9 wherein the protein is produced by recombinant technology.

35

11. A recombinant Eimeria tenella fusion protein immunogen comprising a CheY fusion protein attached by a linking protein which is attached to the amino acid sequence of Claim 9 and microheterogeneous or subunit immunogen forms thereof.

40

12. A purified native Eimeria tenella immunogen protein having an amino acid sequence contained within the amino acid sequence of Claim 9 and any microheterogeneous or subunit forms thereof.

45

13. A method of preparing the protein immunogens of Claim 11 comprising:
 a. obtaining a DNA molecule coding for the protein;
 b. inserting the DNA molecule into an appropriate expression vector;
 c. incorporating the expression vector into an appropriate host cell;
 d. growing the host cell with the expression vector under conditions permitting expression of the DNA and production of the protein; and
 e. recovery of the protein.

50

14. A method of preparing the protein immunogens of Claim 12 comprising:
 a. disrupting Eimeria tenella sporulated oocysts in the presence of protease inhibitors;
 b. contacting the disrupted Eimeria tenella sporulated oocysts with a reducing agent;
 d. carboxymethylation of the reduced solubilized sporulated oocysts material; and
 e. separately recovering the protein immunogen by size exclusion chromatography.

55

15. A method of preparing the protein immunogen of Claim 12 comprising:
 a. disrupting Eimeria tenella sporulated oocysts in the presence of protease inhibitors;
 b. contacting the disrupted Eimeria tenella sporulated oocysts with a reducing agent;

- d. carboxymethylation of the reduced solubilized sporulated oocysts material; and
- e. separately recovering the protein immunogen by immunoaffinity chromatography.

- 5 16. An Eimeria tenella immunogen composition comprising an immunologically effective amount of the immunogen as claimed in any one of Claims 8, 9, 11 or 12 in a physiologically acceptable medium.
- 10 17. An Eimeria tenella immunogen composition comprising an immunologically effective amount of the immunogen of Claim 9 combined with an effective amount of one or more of the immunogens of Group A, Group C, Group H or Group F.
- 15 18. An Eimeria tenella immunogen composition comprising an immunologically effective amount of the immunogen of Claim 12 combined with an effective amount of one or more of the immunogens of Group A, Group C, Group H or Group F.
- 20 19. The use of the immunogen as claimed in any one of Claims 8, 9, 11 or 12, for the preparation of a composition useful for immunizing poultry against Eimeria tenella induced coccidiosis.
- 25 20. A monospecific antibody reactive with the protein immunogen as claimed in any one of Claims 8 to 11.
- 30 21. An Eimeria species Group B protein immunogen comprising an epitope or antigenic determinant corresponding to a Group B Eimeria tenella immunogen encoded by the DNA of Claim 1 which is identified by antibody specific for the Group B Eimeria tenella immunogen.
- 35 22. The Eimeria species Group B immunogen of Claim 21 wherein the species is selected from a group of Eimeria species consisting of E. acervulina, E. mivati, E. mitis, E. preacox, E. hagani, E. necatrix, E. maxima and E. burnetti.
- 40 23. An Eimeria Group B protein immunogen comprising an epitope or antigenic determinant corresponding to the amino acid sequence of Claim 9 and microheterogeneous or subunit immunogen forms thereof and which is selected from a group of Eimeria species consisting of E. acervulina, E. mivati, E. mitis, E. preacox, E. hagani, E. necatrix, E. maxima and E. burnetti.
- 45 24. An Eimeria immunogen composition comprising an immunologically effective amount of one or more of the Group B immunogens of Claim 23.
- 50 25. The use of the immunogen of Claim 23 for the preparation of a composition useful for immunizing poultry against Eimeria induced coccidiosis.

Claims for the following Contracting State : ES

- 55 1. A method of preparing a recombinant DNA molecule comprising a nucleotide sequence corresponding to all or a portion of the following nucleotide sequence:

T CTC GCC CCA ACT TTT TCC CCC GCG CTC CGC AGC AGC AGC AGC AGC AGC AGC AGC AGC
 AGC AAA ATG GCA GAC CTC TTC AGC GGA CTC GTG GGC GGC GTC GTC GGC GCT GTT GCT
 5 GCA GCA GAT TTG CCT GCG GAG GGC GAG AGG GCC CCC CGC CCC GCC CCC GGC ACT GCC
 TGG ACT TGC TGC TGC AGC AAA CTG CAA GAA GGG GCC CGC GAG CTG GAG GGT TTT GTG
 CAG CAG CTG AGT TTT GTT GCA GGG AAG CTG GCC TGC TGC CTG CGG GTG GGG GCG GAG
 CAG CTG GCG CGC TGC GCT GCG GAG GGG CGG CTG CCC AGC AGC AGC AGC AGC AGC AGC
 10 TGC TGC GCG CTG CTG CAG CTC GAG AAG CAG GAC CTC GAG CAG AGC CTC GAG GCC GGC
 AAG CAG GGC GCG GAG TGC CTC TTG AGG AGC AGC AAA CTG GCC CTC GAG GCC CTC CTC
 GAG GGG GCC CGC GTT GCA GCA ACG CGG GGT TTG CTG CTG GTC GAG AGC AGC AAA GAC
 15 ACG GTG CTG CGC AGC ATT CCC CAC ACC CAG GAG AAG CTG GCC CAG GCC TAC AGT TCT
 TTC CTG CGG GGC TAC CAG GGG GCA GCA GCG GGG AGG TCT CTG GGC TAC GGG GCC CCT
 GCT GCT GCT TAC GGC CAG CAG CAG CAG CCC AGC AGC TAC GGG GCG CCC CCC GCC TCC
 AGC CAG CAG CCC TCC GGC TTC TTC TGG TAG CCC TGC AGC AGC AGC AGC AGC AGC AGC
 20 AGC AGC AGC AGC GCG GGC GGC AGC CGC GGC GGG GCC GGG GCG CCG CTG CAG CAA CAG
 CAG CAG CCG nnn CGG CTA GCG CCG CGG AGC ACT CGC AGG GAA CTC CAC AGG CAG CGG
 GAG AGC AGC AGG GAC GAG AAG CAG GTC ATG TAG CGC AGG CAG CAG CGC CAG CTG CAG
 25 CAG CAG CAG CAG CAG CAG CAG CAG CAG CAG CAG CTC CTG CAC CG

which encodes a Group B immunogen of *Eimeria tenella*, which comprises obtaining said DNA molecule by recombinant DNA technology, and combining same with a cloning vector,

- 30 2. A method according to claim 1, wherein said DNA sequence is directly preceded by and operably attached to a second DNA nucleotide sequence resulting in a fused DNA molecule.
3. A method according to claim 2 wherein said second DNA sequence expresses a CheY protein and the product of linking nucleotides.
- 35 4. A method of preparing a recombinant DNA molecule comprising a nucleotide sequence capable of being expressed as at least one protein displaying the immunogenicity of a Group B immunogen of *Eimeria tenella* which sequence is:

40

45

50

55

T CTC GCC CCA ACT TTT TCC CCC GCG CTC CGC AGC AGC AGC AGC AGC AGC AGC AGC AGC
 AGC AAA ATG GCA GAC CTC TTC AGC GGA CTC GTG GGC GGC GTC GTC GGC GCT GTT GCT
 5 GCA GCA GAT TTG CCT GCG GAG GGC GAG AGG GCC CCC CGC CCC GCC CCC GGC ACT GCC
 TGG ACT TGC TGC TGC AGC AAA CTG CAA GAA GGG GCC CGC GAG CTG GAG GGT TTT GTG
 CAG CAG CTG AGT TTT GTT GCA GGG AAG CTG GCC TGC TGC CTG CGG GTG GGG GCG GAG
 CAG CTG GCG CGC TGC GCT GCG GAG GGG CGG CTG CCC AGC AGC AGC AGC AGC AGC AGC
 10 TGC TGC GCG CTG CTG CAG CTC GAG AAG CAG GAC CTC GAG CAG AGC CTC GAG GCC GGC
 AAG CAG GGC GCG GAG TGC CTC TTG AGG AGC AGC AAA CTG GCC CTC GAG GCC CTC CTC
 GAG GGG GCC CGC GTT GCA GCA ACG CGG GGT TTG CTG CTG GTC GAG AGC AGC AAA GAC
 15 ACG GTG CTG CGC AGC ATT CCC CAC ACC CAG GAG AAG CTG GCC CAG GCC TAC AGT TCT
 TTC CTG CGG GGC TAC CAG GGG GCA GCA GCG GGG AGG TCT CTG GGC TAC GGG GCC CCT
 GCT GCT GCT TAC GGC CAG CAG CAG CAG CCC AGC AGC TAC GGG GCG CCC CCC GCC TCC
 AGC CAG CAG CCC TCC GGC TTC TTC TGG TAG CCC TGC AGC AGC AGC AGC AGC AGC AGC
 20 AGC AGC AGC AGC GCG GGC GGC AGC CGC GGC GGG GCC GGG GCG CCG CTG CAG CAA CAG
 CAG CAG CCG nnn CGG CTA GCG CCG CGG AGC ACT CGC AGG GAA CTC CAC AGG CAG CGG
 GAG AGC AGC AGG GAC GAG AAG CAG GTC ATG TAG CGC AGG CAG CAG CGC CAG CTG CAG
 25 CAG CAG CAG CAG CAG CAG CAG CAG CAG CAG CTC CTG CAC CG.

which comprises obtaining said DNA molecule by recombinant DNA technology, and combining same with a cloning vector.

- 30 5. A method according to claim 4, wherein the DNA sequence is directly preceded by and operably attached to a second DNA nucleotide sequence resulting in a fused DNA molecule.
6. A method according to claim 5, wherein said second DNA sequence expresses a CheY protein and the product of linking nucleotides.
- 35 7. A method according to claims 1, 2 or 5, wherein the DNA sequence is contained in an expression vector capable of expressing all or a portion of the immunogen DNA when incorporated in a host cell.
- 40 8. A method of preparing a recombinant Eimeria tenella protein immunogen which comprises expressing the Group B DNA molecule of claim 1 and any microheterogeneous or subunit immunogen forms thereof.
- 45 9. A method of preparing a recombinant Eimeria tenella fusion protein immunogen comprising a CheY fusion protein attached by a linking protein which is attached to an aminoacid sequence contained in a Group B Eimeria tenella protein immunogen, and microheterogeneous or subunit immunogen forms thereof, said method is characterised by:
 - a. obtaining a DNA molecule coding for the protein;
 - b. inserting the DNA molecule into an appropriate expression vector;
 - c. incorporating the expression vector into an appropriate host cell;
 - 50 d. growing the host cell with the expression vector under conditions permitting expression of the DNA and production of the protein; and
 - e. recovery of the protein.
- 55 10. A method according to claim 9, wherein the aminoacid sequence contained in a Group B Eimeria tenella protein immunogen comprises at least the following amino acid sequence:

10

Leu-Ala-Pro-Thr-Phe-Ser-Pro-Ala-Leu-Arg-Ser-Ser-Ser-Ser-Ser-Ser-Ser-

5

20

30

Ser-Ser-Lys-Met-Ala-Asp-Leu-Phe-Ser-Gly-Leu-Val-Gly-Gly-Val-Val-Gly-Ala-

10

40

50

Val-Ala-Ala-Ala-Asp-Leu-Pro-Ala-Glu-Gly-Glu-Arg-Ala-Pro-Arg-Pro-Ala-Pro-

15

60

70

Gly-Thr-Ala-Trp-Thr-Cys-Cys-Cys-Ser-Lys-Leu-Gln-Glu-Gly-Ala-Arg-Glu-

20

80

Leu-Glu-Gly-Phe-Val-Gln-Gln-Leu-Ser-Phe-Val-Ala-Gly-Lys-Leu-Ala-Cys-Cys-

90

100

25

Leu-Arg-Val-Gly-Ala-Glu-Gln-Leu-Ala-Arg-Cys-Ala-Ala-Glu-Gly-Arg-Leu-Pro-

110

120

30

Ser-Ser-Ser-Ser-Ser-Ser-Ser-Ser-Cys-Cys-Ala-Leu-Leu-Gln-Leu-Glu-Lys-Gln-Asp-

130

140

35

Leu-Glu-Gln-Ser-Leu-Glu-Ala-Gly-Lys-Gln-Gly-Ala-Glu-Cys-Leu-Leu-Arg-Ser-

150

160

40

Ser-Lys-Leu-Ala-Leu-Glu-Ala-Leu-Leu-Glu-Gly-Ala-Arg-Val-Ala-Ala-Thr-Agr-

170

45

Gly-Leu-Leu-Leu-Val-Glu-Ser-Ser-Lys-Asp-Thr-Val-Leu-Arg-Ser-Ile-Pro-His-

50

55

180
 Thr-Gln-Glu-Lys-Leu-Ala-Gln-Ala-Tyr-Ser-Ser-Phe-Leu-Arg-Gly-Tyr-Gln-Gly-
 190
 200
 Ala-Ala-Ala-Gly-Arg-Ser-Leu-Gly-Tyr-Gly-Ala-Pro-Ala-Ala-Ala-Tyr-Gly-Gln-
 210
 220
 Gln-Gln-Gln-Pro-Ser-Ser-Tyr-Gly-Ala-Pro-Pro-Ala-Ser-Ser-Gln-Gln-Pro-Ser-
 230
 Gly-Phe-Phe-Trp

and microheterogeneous or subunit immunogen forms thereof.

11. A method according to claim 9, wherein the protein of the Group B Eimeria tenella immunogen is produced by recombinant technology.
12. A method of preparing a purified native Eimeria tenella immunogen protein having an amino acid sequence contained within the aminoacid sequence of a Group B Eimeria tenella and any microheterogeneous or subunit forms thereof, which comprises:
 - a. disrupting Eimeria tenella sporulated oocysts in the presence of protease inhibitors;
 - b. contacting the disrupted Eimeria tenella sporulated oocysts with a reducing agent;
 - c. carboxymethylation of the reduced solubilized sporulated oocysts material; and
 - d. separately recovering the protein immunogen by size exclusion chromatography.
13. A method according to claim 12, wherein the aminoacid sequence comprises at least the following amino acid sequence:

10

Leu-Ala-Pro-Thr-Phe-Ser-Pro-Ala-Leu-Arg-Ser-Ser-Ser-Ser-Ser-Ser-Ser-

5

20

30

Ser-Ser-Lys-Met-Ala-Asp-Leu-Phe-Ser-Gly-Leu-Val-Gly-Gly-Val-Val-Gly-Ala-

10

40

50

Val-Ala-Ala-Ala-Asp-Leu-Pro-Ala-Glu-Gly-Glu-Arg-Ala-Pro-Arg-Pro-Ala-Pro-

15

60

70

Gly-Thr-Ala-Trp-Thr-Cys-Cys-Cys-Ser-Lys-Leu-Gln-Glu-Gly-Ala-Arg-Glu-

20

80

Leu-Glu-Gly-Phe-Val-Gln-Gln-Leu-Ser-Phe-Val-Ala-Gly-Lys-Leu-Ala-Cys-Cys-

90

100

25

Leu-Arg-Val-Gly-Ala-Glu-Gln-Leu-Ala-Arg-Cys-Ala-Ala-Glu-Gly-Arg-Leu-Pro-

110

120

30

Ser-Ser-Ser-Ser-Ser-Ser-Ser-Cys-Cys-Ala-Leu-Leu-Gln-Leu-Glu-Lys-Gln-Asp-

130

140

35

Leu-Glu-Gln-Ser-Leu-Glu-Ala-Gly-Lys-Gln-Gly-Ala-Glu-Cys-Leu-Leu-Arg-Ser-

150

160

40

Ser-Lys-Leu-Ala-Leu-Glu-Ala-Leu-Leu-Glu-Gly-Ala-Arg-Val-Ala-Ala-Thr-Agr-

170

45

Gly-Leu-Leu-Leu-Val-Glu-Ser-Ser-Lys-Asp-Thr-Val-Leu-Arg-Ser-Ile-Pro-His-

50

55

180

190

Thr-Gln-Glu-Lys-Leu-Ala-Gln-Ala-Tyr-Ser-Ser-Phe-Leu-Arg-Gly-Tyr-Gln-Gly-

5

200

210

Ala-Ala-Ala-Gly-Arg-Ser-Leu-Gly-Tyr-Gly-Ala-Pro-Ala-Ala-Ala-Tyr-Gly-Gln-

10

220

230

Gln-Gln-Gln-Pro-Ser-Ser-Tyr-Gly-Ala-Pro-Pro-Ala-Ser-Ser-Gln-Gln-Pro-Ser-

15

Gly-Phe-Phe-Trp

and microheterogeneous or subunit immunogen forms thereof.

20

14. A method of preparing a purified native Eimeria tenella immunogen protein having an aminoacid sequence contained within the aminoacid sequence mentioned in claim 13 and any heterogeneous or subunit forms thereof, which comprises:

25

- a. disrupting Eimeria tenella sporulated oocysts in the presence of protease inhibitors;
- b. contacting the disrupted Eimeria tenella sporulated oocysts with a reducing agent;
- c. carboxymethylation of the reduced solubilized sporulated oocysts material; and
- d. separately recovering the protein immunogen by immunoaffinity chromatography.

30

15. A method according to claims 9, 12 or 14 wherein the Eimeria species Group B protein immunogen comprises an epitope or antigenic determinant corresponding to a Group B Eimeria tenella immunogen encoded by the DNA of Claim 1 which is identified by antibody specific for the Group B Eimeria tenella immunogen.

35

16. A method according to claim 15, wherein the species is selected from a group of Eimeria tenella species consisting of E. acervulina, E. mivati, E. mitis, E. precox, E. hagani, E. necatrix, E. maxima and burnetti.

40

17. A method according to claims 9, 12 or 14 wherein the immunogen comprises an epitope or antigenic determinant corresponding to the amino acid sequence of claim 13 and microheterogeneous or subunit immunogen forms thereof and which is selected from a group of Eimeria species consisting of E. acervulina, E. mivati, E. mitis, E. precox, E. hagani, E. necatrix, E. maxima and E. burnetti.

Patentansprüche

45

Patentansprüche für folgende Vertragsstaaten : AT, BE, CH, LI, DE, FR, GB, IT, LU, NL

1. Rekombinantes DNA-Molekül, umfassend eine Nucleotidsequenz, die vollständig oder teilweise der folgenden Nucleotidsequenz entspricht:

50

55

T CTC GCC CCA ACT TTT TCC CCC GCG CTC CGC AGC AGC AGC AGC AGC AGC AGC AGC AGC
 AGC AAA ATG GCA GAC CTC TTC AGC GGA CTC GTG GGC GGC GTC GTC GGC GCT GTT GCT
 GCA GCA GAT TTG CCT GCG GAG GGC GAG AGG GCC CCC CGC CCC GCC CCC GGC ACT GCC
 TGG ACT TGC TGC TGC AGC AAA CTG CAA GAA GGG GCC CGC GAG CTG GAG GGT TTT GTG
 CAG CAG CTG AGT TTT GTT GCA GGG AAG CTG GCC TGC TGC CTG CGG GTG GGG GCG GAG
 CAG CTG GCG CGC TGC GCT GCG GAG GGG CGG CTG CCC AGC AGC AGC AGC AGC AGC AGC
 TGC TGC GCG CTG CTG CAG CTC GAG AAG CAG GAC CTC GAG CAG AGC CTC GAG GCC GGC
 AAG CAG GGC GCG GAG TGC CTC TTG AGG AGC AGC AAA CTG GCC CTC GAG GCC CTC CTC
 GAG GGG GCC CGC GTT GCA GCA ACG CGG GGT TTG CTG CTG GTC GAG AGC AGC AAA GAC
 ACG GTG CTG CGC AGC ATT CCC CAC ACC CAG GAG AAG CTG GCC CAG GCC TAC AGT TCT
 TTC CTG CGG GGC TAC CAG GGG GCA GCA GCG GGG AGG TCT CTG GGC TAC GGG GCC CCT
 GCT GCT GCT TAC GGC CAG CAG CAG CAG CCC AGC AGC TAC GGG GCG CCC CCC GCC TCC
 AGC CAG CAG CCC TCC GGC TTC TTC TGG TAG CCC TGC AGC AGC AGC AGC AGC AGC AGC
 AGC AGC AGC AGC GCG GGC GGC AGC CGC GGC GGG GCC GGG GCG CCG CTG CAG CAA CAG
 CAG CAG CCG AAA CGG CTA GCG CCG CGG AGC ACT CGC AGG GAA CTC CAC AGG CAG CGG
 GAG AGC AGC AGG GAC GAG AAG CAG GTC ATG TAG CGC AGG CAG CAG CGC CAG CTG CAG
 CAG CAG CAG CAG CAG CAG CAG CAG CAG CAG CAG CTC CTG CAC CG

welches für ein Eimeria tenella-Immunogen der Gruppe B kodiert.

2. Rekombinantes DNA-Molekül nach Anspruch 1, worin der DNA-Sequenz direkt eine Zweite DNA-Nucleotidsequenz vorangeht und die DNA-Sequenz handhabbar an die zweite DNA-Nucleotidsequenz angeknüpft ist, wodurch sich ein DNA-Fusionsmolekül ergibt.
3. Zweite DNA-Sequenz nach Anspruch 2, worin die DNA-Sequenz ein CHeY-Protein und das Produkt der Verknüpfungsnucleotide exprimiert.
4. Rekombinantes DNA-Molekül, umfassend eine Nucleotidsequenz, die in der Lage ist, als mindestens ein Protein exprimiert zu werden, welches die Immunogenität eines Eimeria tenella-Immunogens der Gruppe B aufweist, dessen Sequenz folgendermaßen lautet:

T CTC GCC CCA ACT TTT TCC CCC GCG CTC CGC AGC AGC AGC AGC AGC AGC AGC AGC AGC
 AGC AAA ATG GCA GAC CTC TTC AGC GGA CTC GTG GGC GGC GTC GTC GGC GCT GTT GCT
 GCA GCA GAT TTG CCT GCG GAG GGC GAG AGG GCC CCC CGC CCC GCC CCC GGC ACT GCC
 5 TGG ACT TGC TGC TGC AGC AAA CTG CAA GAA GGG GCC CGC GAG CTG GAG GGT TTT GTG
 CAG CAG CTG AGT TTT GTT GCA GGG AAG CTG GCC TGC TGC CTG CGG GTG GGG GCG GAG
 CAG CTG GCG CGC TGC GCT GCG GAG GGG CGG CTG CCC AGC AGC AGC AGC AGC AGC AGC
 TGC TGC GCG CTG CTG CAG CTC GAG AAG CAG GAC CTC GAG CAG AGC CTC GAG GCC GGC
 10 AAG CAG GGC GCG GAG TGC CTC TTG AGG AGC AGC AAA CTG GCC CTC GAG GCC CTC CTC
 GAG GGG GCC CGC GTT GCA GCA ACG CGG GGT TTG CTG CTG GTC GAG AGC AGC AAA GAC
 ACG GTG CTG CGC AGC ATT CCC CAC ACC CAG GAG AAG CTG GCC CAG GCC TAC AGT TCT
 15 TTC CTG CGG GGC TAC CAG GGG GCA GCA GCG GGG AGG TCT CTG GGC TAC GGG GCC CCT
 GCT GCT GCT TAC GGC CAG CAG CAG CAG CCC AGC AGC TAC GGG GCG CCC CCC GCC TCC
 AGC CAG CAG CCC TCC GGC TTC TTC TGG TAG CCC TGC AGC AGC AGC AGC AGC AGC AGC
 AGC AGC AGC AGC GCG GGC GGC AGC CGC GGC GGG GCC GGG GCG CCG CTG CAG CAA CAG
 20 CAG CAG CCG nnn CGG CTA GCG CCG CGG AGC ACT CGC AGG GAA CTC CAC AGG CAG CGG
 GAG AGC AGC AGG GAC GAG AAG CAG GTC ATG TAG CGC AGG CAG CAG CGC CAG CTG CAG
 CAG CAG CAG CAG CAG CAG CAG CAG CAG CAG CAG CTC CTG CAC CG

- 25
5. Rekombinantes DNA-Molekül nach Anspruch 4, worin der DNA-Sequenz direkt eine zweite DNA-Nucleotidsequenz vorangeht und die DNA-Sequenz handhabbar an die zweite DNA-Nucleotidsequenz angeknüpft ist, wodurch sich ein DNA-Fusionsmolekül ergibt.
- 30
6. Zweite DNA-Sequenz nach Anspruch 5, worin die DNA-Sequenz ein ChEY-Protein und das Produkt der Verknüpfungsnucleotide exprimiert.
- 35
7. Rekombinantes DNA-Molekül nach einem der Ansprüche 1, 2 oder 5, worin die DNA-Sequenz in einem Expressionsvektor enthalten ist, der in der Lage ist, die immunogene DNA ganz oder teilweise zu exprimieren, wenn er in eine Wirtszelle eingebaut wird.
- 40
8. Rekombinantes *Eimeria tenella*-Proteinimmunogen, umfassend eine Aminosäuresequenz, die durch das DNA-Molekül der Gruppe B nach Anspruch 1 exprimiert wird und irgendwelche mikroheterogenen oder immunogenen Untereinheit-Formen davon.
- 45
9. *Eimeria tenella*-Proteinimmunogen der Gruppe B, umfassend mindestens die folgende Aminosäuresequenz:
- 50
- 55

10

Leu-Ala-Pro-Thr-Phe-Ser-Pro-Ala-Leu-Arg-Ser-Ser-Ser-Ser-Ser-Ser-Ser-

5

20

30

Ser-Ser-Lys-Met-Ala-Asp-Leu-Phe-Ser-Gly-Leu-Val-Gly-Gly-Val-Val-Gly-Ala-

10

40

50

Val-Ala-Ala-Ala-Asp-Leu-Pro-Ala-Glu-Gly-Glu-Arg-Ala-Pro-Arg-Pro-Ala-Pro-

15

60

70

Gly-Thr-Ala-Trp-Thr-Cys-Cys-Cys-Ser-Lys-Leu-Gln-Glu-Gly-Ala-Arg-Glu-

20

80

Leu-Glu-Gly-Phe-Val-Gln-Gln-Leu-Ser-Phe-Val-Ala-Gly-Lys-Leu-Ala-Cys-Cys-

25

90

100

Leu-Arg-Val-Gly-Ala-Glu-Gln-Leu-Ala-Arg-Cys-Ala-Ala-Glu-Gly-Arg-Leu-Pro-

30

110

120

Ser-Ser-Ser-Ser-Ser-Ser-Ser-Ser-Cys-Cys-Ala-Leu-Leu-Gln-Leu-Glu-Lys-Gln-Asp-

35

130

140

Leu-Glu-Gln-Ser-Leu-Glu-Ala-Gly-Lys-Gln-Gly-Ala-Glu-Cys-Leu-Leu-Arg-Ser-

40

150

160

Ser-Lys-Leu-Ala-Leu-Glu-Ala-Leu-Leu-Glu-Gly-Ala-Arg-Val-Ala-Ala-Thr-Agr-

45

170

Gly-Leu-Leu-Leu-Val-Glu-Ser-Ser-Lys-Asp-Thr-Val-Leu-Arg-Ser-Ile-Pro-His-

180

190

Thr-Gln-Glu-Lys-Leu-Ala-Gln-Ala-Tyr-Ser-Ser-Phe-Leu-Arg-Gly-Tyr-Gln-Gly-

50

55

200

210

Ala-Ala-Ala-Gly-Arg-Ser-Leu-Gly-Tyr-Gly-Ala-Pro-Ala-Ala-Ala-Tyr-Gly-Gln-

5

220

230

Gln-Gln-Gln-Pro-Ser-Ser-Tyr-Gly-Ala-Pro-Pro-Ala-Ser-Ser-Gln-Gln-Pro-Ser-

10

Gly-Phe-Phe-Trp .

und die mikroheterogenen oder immunogenen Untereinheit-Formen davon.

15 10. *Eimeria tenella*-Immunogen der Gruppe B nach Anspruch 9, worin das Protein durch Rekombinationstechnik hergestellt wird.

11. Rekombinantes *Eimeria tenella*-Fusionsprotein-Immunogen, umfassend ein CheY-Fusionsprotein, das von einem Verknüpfungsprotein angeknüpft wird, das an die Aminosäuresequenz nach Anspruch 9 angeknüpft wird und die mikroheterogenen oder immunogenen Untereinheit-Formen davon.

12. Gereinigtes natives *Eimeria tenella*-Immunogenprotein mit einer Aminosäuresequenz, die in der Aminosäuresequenz nach Anspruch 9 enthalten ist und irgendeine mikroheterogene oder Form der Untereinheit davon.

25

13. Verfahren zur Herstellung von Proteinimmunogen nach Anspruch 11 umfassend:

- a. Erhalten eines DNA-Moleküls, welches für das Protein kodiert;
- b. Inserieren des DNA-Moleküls in einen geeigneten Expressionsvektor;
- c. Einbauen des Expressionsvektors in eine geeignete Wirtszelle;
- d. Züchten der Wirtszelle mit dem Expressionsvektor unter Bedingungen, die die Expression der DNA und Herstellung der Proteine erlauben; und
- e. Gewinnen des Proteins.

30

14. Verfahren zur Herstellung von Proteinimmunogenen nach Anspruch 12, umfassend:

- a. Aufbrechen von sporulierten *Eimeria tenella*-Oozysten in Gegenwart von Proteaseinhibitoren;
- b. Kontaktieren der aufgebrochenen sporulierten *Eimeria tenella*-Oozysten mit einem Reduktionsmittel;
- d. Carboxymethylieren des reduzierten, solubilisierten, sporulierten Oozystenmaterials; und
- e. getrennt Gewinnen des Proteinimmunogens durch Größenausschlußchromatographie.

40

15. Verfahren zur Herstellung des Proteinimmunogens nach Anspruch 12, umfassend:

- a. Aufbrechen von sporulierten *Eimeria tenella*-Oozysten in Gegenwart von Proteaseinhibitoren;
- b. Kontaktieren der aufgebrochenen sporulierten *Eimeria tenella*-Oozysten mit einem Reduktionsmittel;
- d. Carboxymethylieren des reduzierten, solubilisierten, sporulierten Oozystenmaterials; und
- e. getrennt Gewinnen des Proteinimmunogens durch Größenausschlußchromatographie.

45

16. *Eimeria tenella*-Immunogenpräparat, umfassend eine immunologisch wirksame Menge des Immunogens nach einem der Ansprüche 8, 9, 11 oder 12 in einem physiologisch verträglichen Medium.

50

17. *Eimeria tenella*-Immunogenpräparat, umfassend eine immunologisch wirksame Menge des Immunogens nach Anspruch 9 in Verbindung mit einer wirksamen Menge eines oder mehrerer Immunogene der Gruppen A, C, H oder F.

18. *Eimeria tenella*-Immunogenpräparat, umfassend eine immunologisch wirksame Menge des Immunogens nach Anspruch 12 in Verbindung mit einer wirksamen Menge eines oder mehrerer Immunogene der Gruppen A, C, H oder F.

55

19. Verwendung des Immunogens nach einem der Ansprüche 8, 9, 11 oder 12 zur Herstellung eines Präparates, welches zur Immunisierung von Geflügel gegen eine durch Eimeria tenella hervorgerufene Kokzidiose geeignet ist.
20. Monospezifischer Antikörper, der mit dem Proteinimmunogen nach einem der Ansprüche 8 bis 11 reaktiv ist.
21. Proteinimmunogen der Gruppe B der Spezies Eimeria, umfassend eine epitope oder antigenische Determinante, die einem Eimeria tenella-Immunogen der Gruppe B entspricht, das von der DNA nach Anspruch 1 codiert worden ist, die durch einen für das Eimeria tenella-Immunogen der Gruppe B spezifischen Antikörper identifiziert wird.
22. Immunogen der Gruppe B der Spezies Eimeria nach Anspruch 1, worin die Spezies ausgewählt ist aus einer Gruppe von Eimeria-Spezies, bestehend aus E. acervulina, E. mivati, E. mitis, E. preacox, E. hagani, E. necatrix, E. maxima und E. burnetti.
23. Eimeria-Proteinimmunogen der Gruppe B, umfassend eine epitope oder antigenische Determinante, die der Aminosäuresequenz nach Anspruch 9 entspricht und die mikroheterogenen oder immunogenen Formen der Unterheit davon, das ausgewählt ist aus einer Gruppe von Eimeria-Spezies, bestehend aus E. acervulina, E. mivati, E. mitis, E. preacox, E. hagani, E. necatrix, E. maxima und E. burnetti.
24. Eimeria-Immunogenpräparat, umfassend eine immunologisch wirksame Menge eines oder mehrerer Immunogene der Gruppe B nach Anspruch 23.
25. Verwendung des Immunogens nach Anspruch 23 zur Herstellung eines Präparates, das zur Immunisierung von Geflügel gegen eine durch Eimeria hervorgerufene Kokzidiose geeignet ist.

Patentansprüche für folgenden Vertragsstaat : ES

1. Verfahren zur Herstellung eines rekombinanten DNA-Moleküls, umfassend eine Nucleotidsequenz, die ganz oder teilweise der folgenden Nucleotidsequenz entspricht:

```

T CTC GCC CCA ACT TTT TCC CCC GCG CTC CGC AGC AGC AGC AGC AGC AGC AGC AGC AGC
AGC AAA ATG GCA GAC CTC TTC AGC GGA CTC GTG GGC GGC GTC GTC GGC GCT GTT GCT
GCA GCA GAT TTG CCT GCG GAG GGC GAG AGG GCC CCC CGC CCC GCC CCC GGC ACT GCC
TGG ACT TGC TGC TGC AGC AAA CTG CAA GAA GGG GCC CGC GAG CTG GAG GGT TTT GTG
CAG CAG CTG AGT TTT GTT GCA GGG AAG CTG GCC TGC TGC CTG CGG GTG GGG GCG GAG
CAG CTG GCG CGC TGC GCT GCG GAG GGG CGG CTG CCC AGC AGC AGC AGC AGC AGC AGC
TGC TGC GCG CTG CTG CAG CTC GAG AAG CAG GAC CTC GAG CAG AGC CTC GAG GCC GGC
AAG CAG GGC GCG GAG TGC CTC TTG AGG AGC AGC AAA CTG GCC CTC GAG GCC CTC CTC
GAG GGG GCC CGC GTT GCA GCA ACG CGG GGT TTG CTG CTG GTC GAG AGC AGC AAA GAC
ACG GTG CTG CGC AGC ATT CCC CAC ACC CAG GAG AAG CTG GCC CAG GCC TAC AGT TCT
TTC CTG CGG GGC TAC CAG GGG GCA GCA GCG GGG AGG TCT CTG GGC TAC GGG GCC CCT
GCT GCT GCT TAC GGC CAG CAG CAG CAG CCC AGC AGC TAC GGG GCG CCC CCC GCC TCC
AGC CAG CAG CCC TCC GGC TTC TTC TGG TAG CCC TGC AGC AGC AGC AGC AGC AGC AGC
AGC AGC AGC AGC GCG GGC GGC AGC CGC GGC GGG GCC GGG GCG CCG CTG CAG CAA CAG
CAG CAG CCG nnn CGG CTA GCG CCG CGG AGC ACT CGC AGG GAA CTC CAC AGG CAG CGG
GAG AGC AGC AGG GAC GAG AAG CAG GTC ATG TAG CGC AGG CAG CAG CGC CAG CTG CAG
CAG CAG CAG CAG CAG CAG CAG CAG CAG CAG CAG CAG CTC CTG CAC CG.

```

das für ein Immunogen der Gruppe B von Eimeria tenella codiert, das umfaßt Erhalten des genannten

DNA-Moleküls durch DNA-Rekombinationstechnik und Kombinieren desselben mit einem Klonierungsvektor.

2. Verfahren nach Anspruch 1, bei dem der genannten DNA-Sequenz direkt eine zweite DNA-Nucleotidsequenz vorangeht und die DNA-Sequenz handhabbar an die zweite DNA-Nucleotidsequenz angeknüpft ist, wodurch sich ein DNA-Fusionsmolekül ergibt.
3. Verfahren nach Anspruch 2, bei dem die genannte zweite DNA-Sequenz ein CheY-Protein und das Produkt von Verknüpfungsnucleotiden exprimiert.
4. Verfahren zur Herstellung eines rekombinanten DNA-Moleküls, das eine Nucleotidsequenz umfaßt, die in der Lage ist, als mindestens ein Protein exprimiert zu werden, welches die Immunogenität eines Immunogens der Gruppe B von Eimeria tenella aufweist, dessen Sequenz folgendermaßen lautet:

```

T CTC GCC CCA ACT TTT TCC CCC GCG CTC CGC AGC AGC AGC AGC AGC AGC AGC AGC AGC
AGC AAA ATG GCA GAC CTC TTC AGC GGA CTC GTG GGC GGC GTC GTC GGC GCT GTT GCT
GCA GCA GAT TTG CCT GCG GAG GGC GAG AGG GCC CCC CGC CCC GCC CCC GGC ACT GCC
TGG ACT TGC TGC TGC AGC AAA CTG CAA GAA GGG GCC CGC GAG CTG GAG GGT TTT GTG
CAG CAG CTG AGT TTT GTT GCA GGG AAG CTG GCC TGC TGC CTG CGG GTG GGG GCG GAG
CAG CTG GCG CGC TGC GCT GCG GAG GGG CGG CTG CCC AGC AGC AGC AGC AGC AGC AGC
TGC TGC GCG CTG CTG CAG CTC GAG AAG CAG GAC CTC GAG CAG AGC CTC GAG GCC GGC
AAG CAG GGC GCG GAG TGC CTC TTG AGG AGC AGC AAA CTG GCC CTC GAG GCC CTC CTC
GAG GGG GCC CGC GTT GCA GCA ACG CGG GGT TTG CTG CTG GTC GAG AGC AGC AAA GAC
ACG GTG CTG CGC AGC ATT CCC CAC ACC CAG GAG AAG CTG GCC CAG GCC TAC AGT TCT
TTC CTG CGG GGC TAC CAG GGG GCA GCA GCG GGG AGG TCT CTG GGC TAC GGG GCC CCT
GCT GCT GCT TAC GGC CAG CAG CAG CAG CCC AGC AGC TAC GGG GCG CCC CCC GCC TCC
AGC CAG CAG CCC TCC GGC TTC TTC TGG TAG CCC TGC AGC AGC AGC AGC AGC AGC AGC
AGC AGC AGC AGC GCG GGC GGC AGC CGC GGC GGG GCC GGG GCG CCG CTG CAG CAA CAG
CAG CAG CCG nnn CGG CTA GCG CCG CGG AGC ACT CGC AGG GAA CTC CAC AGG CAG CGG
GAG AGC AGC AGG GAC GAG AAG CAG GTC ATG TAG CGC AGG CAG CAG CGC CAG CTG CAG
CAG CAG CAG CAG CAG CAG CAG CAG CAG CAG CTC CTG CAC CG.

```

welches umfaßt Erhalten des genannten DNA-Moleküls durch DNA-Rekombinationstechnik und Kombinieren desselben mit einem Klonierungsvektor.

5. Verfahren nach Anspruch 4, bei dem der DNA-Sequenz direkt eine zweite DNA-Nucleotidsequenz vorangeht und die DNA-Sequenz handhabbar an die zweite DNA-Nucleotidsequenz angeknüpft ist, wodurch sich ein DNA-Fusionsmolekül ergibt.
6. Verfahren nach Anspruch 5, bei dem die genannte zweite DNA-Sequenz ein CheY-Protein und das Produkt der Verknüpfungsnucleotide exprimiert.
7. Verfahren nach den Ansprüchen 1, 2 oder 5, bei dem die DNA-Sequenz in einem Expressionsvektor enthalten ist, der in der Lage ist, die immunogene DNA ganz oder teilweise zu exprimieren, wenn er in eine Wirtszelle eingebaut wird.
8. Verfahren zur Herstellung eines rekombinanten Eimeria tenella-Proteinimmunogen, welches umfaßt Exprimieren des DNA-Moleküls der Gruppe B nach Anspruch 1 und irgendeiner mikroheterogenen oder immunogen Form der Untereinheit davon.

9. Verfahren zur Herstellung eines rekombinanten Fusionsproteinimmunogens aus Eimeria tenella, umfassend ein CheY-Fusionsprotein, das von einem Verknüpfungspotein angeknüpft wird, das an eine Aminosäuresequenz angeknüpft wird, die in einem Eimeria tenella-Proteinimmunogen der Gruppe B enthalten ist und einer mikroheterogenen oder immunogenen Untereinheit-Form davon, wobei das genannte Verfahren gekennzeichnet ist durch:
 - a. Erhalten eines DNA-Moleküls, welches für das Protein kodiert;
 - b. Inserieren des DNA-Moleküls in einen geeigneten Expressionsvektor;
 - c. Einbau des Expressionsvektors in eine geeignete Wirtszelle;
 - d. Züchten der Wirtszelle mit dem Expressionsvektor unter Bedingungen, die die Expression der DNA und Herstellung der Proteine erlauben; und
 - e. Gewinnen des Proteins.
10. Verfahren nach Anspruch 9, bei dem die Aminosäuresequenz, die in einem Eimeria tenella-Proteinimmunogen der Gruppe B enthalten ist, mindestens die folgende Aminosäuresequenz:

10
Leu-Ala-Pro-Thr-Phe-Ser-Pro-Ala-Leu-Arg-Ser-Ser-Ser-Ser-Ser-Ser-Ser-Ser-

20 30
Ser-Ser-Lys-Met-Ala-Asp-Leu-Phe-Ser-Gly-Leu-Val-Gly-Gly-Val-Val-Gly-Ala-

40 50
Val-Ala-Ala-Ala-Asp-Leu-Pro-Ala-Glu-Gly-Glu-Arg-Ala-Pro-Arg-Pro-Ala-Pro-

60 70
Gly-Thr-Ala-Trp-Thr-Cys-Cys-Cys-Ser-Lys-Leu-Gln-Glu-Gly-Ala-Arg-Glu-

80
Leu-Glu-Gly-Phe-Val-Gln-Gln-Leu-Ser-Phe-Val-Ala-Gly-Lys-Leu-Ala-Cys-Cys-

90 100
Leu-Arg-Val-Gly-Ala-Glu-Gln-Leu-Ala-Arg-Cys-Ala-Ala-Glu-Gly-Arg-Leu-Pro-

110 120
Ser-Ser-Ser-Ser-Ser-Ser-Ser-Cys-Cys-Ala-Leu-Leu-Gln-Leu-Glu-Lys-Gln-Asp-

130 140
Leu-Glu-Gln-Ser-Leu-Glu-Ala-Gly-Lys-Gln-Gly-Ala-Glu-Cys-Leu-Leu-Arg-Ser-

150 160
 Ser-Lys-Leu-Ala-Leu-Glu-Ala-Leu-Leu-Glu-Gly-Ala-Arg-Val-Ala-Ala-Thr-Agr-
 5
 170
 Gly-Leu-Leu-Leu-Val-Glu-Ser-Ser-Lys-Asp-Thr-Val-Leu-Arg-Ser-Ile-Pro-His-
 10
 180 190
 Thr-Gln-Glu-Lys-Leu-Ala-Gln-Ala-Tyr-Ser-Ser-Phe-Leu-Arg-Gly-Tyr-Gln-Gly-
 15
 200 210
 Ala-Ala-Ala-Gly-Arg-Ser-Leu-Gly-Tyr-Gly-Ala-Pro-Ala-Ala-Ala-Tyr-Gly-Gln-
 20
 220 230
 Gln-Gln-Gln-Pro-Ser-Ser-Tyr-Gly-Ala-Pro-Pro-Ala-Ser-Ser-Gln-Gln-Pro-Ser-
 25
 Gly-Phe-Phe-Trp

und die mikroheterogenen oder immunogenen Untereinheit-Formen davon umfaßt.

11. Verfahren nach Anspruch 9, bei dem das Protein des Eimeria tenella-Immunogens der Gruppe B durch Rekombinationstechnik hergestellt wird.
12. Verfahren zur Herstellung eines gereinigten nativen Eimeria tenella-Immunogenproteins das eine Aminosäuresequenz in der Eimeria tenella-Aminosäuresequenz der Gruppe B enthält und irgendeiner mikroheterogenen oder Untereinheit-Form davon, welches umfaßt:
 - a. Aufbrechen von sporulierten Eimeria tenella-Oozysten in Gegenwart von Proteaseinhibitoren;
 - b. Kontaktieren der aufgebrochenen, sporulierten Eimeria tenella-Oozysten mit einem Reduktionsmittel
 - d. Carboxymethylieren des reduzierten, solubilisierten, sporulierten Oozystenmaterials; und
 - e. getrennt Gewinnen des Proteinimmunogens durch Größenausschlußchromatographie.
13. Verfahren nach Anspruch 12, bei dem die Aminosäuresequenz mindestens die folgende Aminosäuresequenz

10

Leu-Ala-Pro-Thr-Phe-Ser-Pro-Ala-Leu-Arg-Ser-Ser-Ser-Ser-Ser-Ser-Ser-

20

30

Ser-Ser-Lys-His-Ala-Asp-Leu-Phe-Ser-Gly-Leu-Val-Gly-Gly-Val-Val-Gly-Ala-

40

50

Val-Ala-Ala-Ala-Asp-Leu-Pro-Ala-Glu-Gly-Glu-Arg-Ala-Pro-Arg-Pro-Ala-Pro-

60

70

Gly-Thr-Ala-Trp-Thr-Cys-Cys-Cys-Ser-Lys-Leu-Gln-Glu-Gly-Ala-Arg-Glu-

80

Leu-Glu-Gly-Phe-Val-Gln-Gln-Leu-Ser-Phe-Val-Ala-Gly-Lys-Leu-Ala-Cys-Cys-

90

100

Leu-Arg-Val-Gly-Ala-Glu-Gln-Leu-Ala-Arg-Cys-Ala-Ala-Glu-Gly-Arg-Leu-Pro-

110

120

Ser-Ser-Ser-Ser-Ser-Ser-Ser-Cys-Cys-Ala-Leu-Leu-Gln-Leu-Glu-Lys-Gln-Asp-

130

140

Leu-Glu-Gln-Ser-Leu-Glu-Ala-Gly-Lys-Gln-Gly-Ala-Glu-Cys-Leu-Leu-Arg-Ser-

150

160

Ser-Lys-His-Ala-Ile-Gln-Ala-Leu-Leu-Glu-Gly-Ala-Arg-Val-Ala-Ala-Thr-Arg-

170

Gly-Leu-Leu-Leu-Val-Glu-Ser-Ser-Lys-Asp-Thr-Val-Leu-Arg-Ser-Ile-Pro-His-

180

190

Thr-Gln-Glu-Lys-Leu-Ala-Gln-Ala-Tyr-Ser-Ser-Phe-Leu-Arg-Gly-Tyr-Gln-Gly-

5

200

210

Ala-Ala-Ala-Gly-Arg-Ser-Leu-Gly-Tyr-Gly-Ala-Pro-Ala-Ala-Ala-Tyr-Gly-Gln-

10

220

230

Gln-Gln-Gln-Pro-Ser-Ser-Tyr-Gly-Ala-Pro-Pro-Ala-Ser-Ser-Gln-Gln-Pro-Ser-

15

Gly-Phe-Phe-Trp

und die mikroheterogenen oder immunogenen Untereinheit-Formen davon umfaßt.

20

14. Verfahren zur Herstellung eines gereinigten nativen Eimeria tenella-Immunogenproteins, das in der Aminosäuresequenz nach Anspruch 13 eine Aminosäuresequenz enthält und irgendwelcher heterogenen oder Untereinheit-Formen davon, welches umfaßt:

25

- a. Aufbrechen von sporulierten Eimeria tenella-Oozysten in Gegenwart von Proteaseinhibitoren;
- b. Kontaktieren der aufgebrochenen sporulierten Eimeria tenella-Oozysten mit einem Reduktionsmittel;
- d. Carboxymethylieren des reduzierten, solubilisierten, sporulierten Oozystenmaterials; und
- e. getrennt Gewinnen des Proteinimmunogens durch Größenausschlußchromatographie.

30

15. Verfahren nach den Ansprüchen 9, 12 oder 14, worin das Proteinimmunogen der Gruppe B der Spezies Eimeria umfaßt eine epitope oder antigenische Determinante, die einem durch die DNA nach Anspruch 1 kodierten Eimeria tenella-Immunogen der Gruppe B entspricht, welches identifiziert wird durch einen Antikörper, der für das Eimeria tenella-Immunogen der Gruppe B spezifisch ist.

35

16. Verfahren nach Anspruch 15, bei dem die Spezies ausgewählt wird aus einer Gruppe von Eimeria tenella-Spezies, bestehend aus E. acervulina, E. mivati, E. mitis, E. preacox, E. hagani, E. necatrix, E. maxima und burnetti.

40

17. Verfahren nach den Ansprüchen 9, 12 oder 14, bei dem das Immunogen umfaßt eine epitope oder antigenische Determinante, die der Aminosäuresequenz nach Anspruch 13 entspricht und die mikroheterogenen oder immunogenen Untereinheit-Formen davon, welches ausgewählt wird aus einer Gruppe von Eimeria-Spezies, bestehend aus E. acervulina, E. mivati, E. mitis, E. preacox, E. hagani, E. necatrix, E. maxima und burnetti.

Revendications

Revendications pour les Etats contractants suivants : AT, BE, CH, LI, DE, FR, GB, IT, LU, NL

45

1. Molécule d'ADN recombinante, comprenant une séquence nucléotidique correspondant à l'ensemble ou à une partie de la séquence nucléotidique suivante:

50

55

25

qui code pour un immunogène de groupe B d'*Eimeria tenella*.

- 30

35

40

45

50

55

1 CTC GCC CCA ACT TTT TCC CCC GCG CTC CGC AGC AGC AGC AGC AGC AGC AGC AGC AGC
 AGC AAA ATG GCA GAC CTC TTC AGC GGA CTC GTG GGC GGC GTC GTC GGC GCT GTT GCT
 GCA GCA GAT TTG CCT GCG GAG GGC GAG AGG GCC CCC CGC CCC GCC CCC GGC ACT GCC
 5 TGG ACT TGC TGC TGC AGC AAA CTG CAA GAA GGG GCC CGC GAG CTG GAG GGT TTT GTG
 CAG CAG CTG AGT TTT GTT GCA GGG AAG CTG GCC TGC TGC CTG CGG GTG GGG GCG GAG
 CAG CTG GCG CGC TGC GCT GCG GAG GGG CGG CTG CCC AGC AGC AGC AGC AGC AGC AGC
 10 TGC TGC GCG CTG CTG CAG CTC GAG AAG CAG GAC CTC GAG CAG AGC CTC GAG GCC GGC
 AAG CAG GGC GCG GAG TGC CTC TTG AGG AGC AGC AAA CTG GCC CTC GAG GCC CTC CTC
 GAG GGG GCC CGC GTT GCA GCA ACG CGG GGT TTG CTG CTG GTC GAG AGC AGC AAA GAC
 ACG GTG CTG CGC AGC ATT CCC CAC ACC CAG GAG AAG CTG GCC CAG GCC TAC AGT TCT
 15 TTC CTG CGG GGC TAC CAG GGG GCA GCA GCG GGG AGG TCT CTG GGC TAC GGG GCC CCT
 GCT GCT GCT TAC GGC CAG CAG CAG CAG CCC AGC AGC TAC GGG GCG CCC CCC GCC TCC
 AGC CAG CAG CCC TCC GGC TTC TTC TGG TAG CCC TGC AGC AGC AGC AGC AGC AGC AGC
 AGC AGC AGC AGC GCG GGC GGC AGC CGC GGC GGG GCC GGG GCG CCG CTG CAG CAA CAG
 20 CAG CAG CCG nnn CGG CTA GCG CCG CGG AGC ACT CGC AGG GAA CTC CAC AGG CAG CGG
 GAG AGC AGC AGG GAC GAG AAG CAG GTC ATG TAG CGC AGG CAG CAG CGC CAG CTG CAG
 CAG CAG CAG CAG CAG CAG CAG CAG CAG CAG CTC CTG CAC CG.

25

5. Molécule d'ADN recombinante selon la revendication 4, dans laquelle la séquence d'ADN est directement précédée par et fonctionnellement liée à une seconde séquence nucléotidique d'ADN, ce qui donne une molécule d'ADN fusionnée.

30

6. Seconde séquence d'ADN de la revendication 5, dans laquelle la séquence d'ADN exprime une protéine CheY et le produit de nucléotides de liaison.

35

7. Molécule d'ADN recombinante selon l'une quelconque des revendications 1, 2 ou 5, dans laquelle la séquence d'ADN est contenue dans un vecteur d'expression capable d'exprimer l'ensemble ou une partie de l'ADN de l'immunogène, lorsqu'elle est incorporée dans une cellule hôte.

40

8. Immunogène protéique recombinant d'*Eimeria tenella*, comprenant une séquence d'acides aminés exprimée par la molécule d'ADN de groupe B de la revendication 1, et toutes formes immunogènes de type sous-unité ou microhétérogènes de celui-ci.

45

9. Immunogène protéique d'*Eimeria tenella* de groupe B, comprenant au moins la séquence d'acides aminés suivante:

50

55

10

Leu-Ala-Pro-Thr-Phe-Ser-Pro-Ala-Leu-Arg-Ser-Ser-Ser-Ser-Ser-Ser-Ser-

5

20

30

Ser-Ser-Lys-Met-Ala-Asp-Leu-Phe-Ser-Gly-Leu-Val-Gly-Gly-Val-Val-Gly-Ala-

10

40

50

Val-Ala-Ala-Ala-Asp-Leu-Pro-Ala-Glu-Gly-Glu-Arg-Ala-Pro-Arg-Pro-Ala-Pro-

15

60

70

Gly-Thr-Ala-Trp-Thr-Cys-Cys-Cys-Ser-Lys-Leu-Gln-Glu-Gly-Ala-Arg-Glu-

80

20

Leu-Glu-Gly-Phe-Val-Gln-Gln-Leu-Ser-Phe-Val-Ala-Gly-Lys-Leu-Ala-Cys-Cys-

90

100

25

Leu-Arg-Val-Gly-Ala-Glu-Gln-Leu-Ala-Arg-Cys-Ala-Ala-Glu-Gly-Arg-Leu-Pro-

110

120

30

Ser-Ser-Ser-Ser-Ser-Ser-Ser-Cys-Cys-Ala-Leu-Leu-Gln-Leu-Glu-Lys-Gln-Asp-

130

140

Leu-Glu-Gln-Ser-Leu-Glu-Ala-Gly-Lys-Gln-Gly-Ala-Glu-Cys-Leu-Leu-Arg-Ser-

35

150

160

Ser-Lys-Leu-Ala-Leu-Glu-Ala-Leu-Leu-Glu-Gly-Ala-Arg-Val-Ala-Ala-Thr-Agr-

40

170

Gly-Leu-Leu-Leu-Val-Glu-Ser-Ser-Lys-Asp-Thr-Val-Leu-Arg-Ser-Ile-Pro-His-

180

190

45

Thr-Gln-Glu-Lys-Leu-Ala-Gln-Ala-Tyr-Ser-Ser-Phe-Leu-Arg-Gly-Tyr-Gln-Gly-

200

210

50

Ala-Ala-Ala-Gly-Arg-Ser-Leu-Gly-Tyr-Gly-Ala-Pro-Ala-Ala-Ala-Tyr-Gly-Gln-

220

230

55

Gln-Gln-Gln-Pro-Ser-Ser-Tyr-Gly-Ala-Pro-Pro-Ala-Ser-Ser-Gln-Gln-Pro-Ser-

Gly-Phe-Phe-Trp

et formes immunogènes de type sous-unité ou microhétérogènes de celui-ci.

10. Immunogène de groupe B d'*Eimeria tenella* selon la revendication 9, dans lequel la protéine est produite par la technique de l'ADN recombinant.
- 5 11. Immunogène de type protéine de fusion recombinante d'*Eimeria tenella*, comprenant une protéine de fusion CheY liée par une protéine de liaison qui est fixée à la séquence d'acides aminés de la revendication 9, et formes immunogènes de type sous-unité ou microhétérogènes de celui-ci.
- 10 12. Protéine immunogène native purifiée d'*Eimeria tenella*, comportant une séquence d'acides aminés contenue dans la séquence d'acides aminés de la revendication 9, et toutes formes de type sous-unité ou microhétérogènes de celle-ci.
- 15 13. Procédé de production des immunogènes protéiques selon la revendication 11, comprenant:
 - a. l'obtention d'une molécule d'ADN codant pour la protéine;
 - b. l'insertion de la molécule d'ADN dans un vecteur d'expression approprié;
 - c. l'incorporation du vecteur d'expression dans une cellule hôte appropriée;
 - d. la culture de la cellule hôte avec le vecteur d'expression, dans des conditions permettant l'expression de l'ADN et la production de la protéine; et
 - 20 e. la récupération de la protéine.
- 25 14. Procédé de production des immunogènes protéiques de la revendication 12, comprenant:
 - a. la désintégration d'ookystes sporulés d'*Eimeria tenella*, en présence d'inhibiteurs de protéase;
 - b. la mise en contact des ookystes sporulés d'*Eimeria tenella*, désintégrés, avec un agent réducteur;
 - c. la carboxyméthylation de la substance d'ookystes sporulés solubilisée, réduite; et
 - d. la récupération séparément de l'immunogène protéique par chromatographie par exclusion de taille.
- 30 15. Procédé de production de l'immunogène protéique de la revendication 12, comprenant:
 - a. la désintégration d'ookystes d'*Eimeria tenella* sporulés, en présence d'inhibiteurs de protéase;
 - b. la mise en contact des ookystes sporulés d'*Eimeria tenella*, désintégrés, avec un agent réducteur;
 - c. la carboxyméthylation de la substance d'ookystes sporulés, solubilisée, réduite; et
 - 35 d. la récupération séparément de l'immunogène protéique par chromatographie d'immunoaffinité.
- 40 16. Composition d'immunogène d'*Eimeria tenella* comprenant une quantité immunologiquement efficace de l'immunogène selon l'une quelconque des revendications 8, 9, 11 ou 12, dans un milieu physiologiquement acceptable.
- 45 17. Composition d'immunogène d'*Eimeria tenella* comprenant une quantité immunologiquement efficace de l'immunogène de la revendication 9, associée avec une quantité efficace d'un ou plusieurs des immunogènes du groupe A, du groupe C, du groupe H ou du groupe F.
- 50 18. Composition d'immunogène d'*Eimeria tenella* comprenant une quantité immunologiquement efficace de l'immunogène de la revendication 12, associée avec une quantité efficace d'un ou plusieurs des immunogènes du groupe A, du groupe C, du groupe H ou du groupe F.
- 55 19. Utilisation de l'immunogène selon l'une quelconque des revendications 8, 9, 11 ou 12, pour la préparation d'une composition utilisable pour l'immunisation de volailles contre la coccidiose provoquée par *Eimeria tenella*.
20. Anticorps monospécifique réactif avec l'immunogène protéique selon l'une quelconque des revendications 8 à 11.
21. Immunogène protéique de groupe B d'une espèce appartenant au genre *Eimeria*, comprenant un épitope ou déterminant antigénique correspondant à un immunogène d'*Eimeria tenella* de groupe B, codé par l'ADN de la revendication 1, qui est identifié par un anticorps spécifique de l'immunogène

d'*Eimeria tenella* de groupe B.

22. Immunogène de groupe B d'une espèce appartenant au genre *Eimeria* de la revendication 21, dans lequel l'espèce est choisie dans un groupe d'espèces appartenant au genre *Eimeria*, comprenant *E. acervulina*, *E. mivati*, *E. mitis*, *E. preacox*, *E. hagani*, *E. necatrix*, *E. maxima* et *E. burnetti*.

23. Immunogène protéique d'*Eimeria* de groupe B, comprenant un épitope ou déterminant antigénique correspondant à la séquence d'acides aminés de la revendication 9 et à des formes immunogènes de type sous-unité ou microhétérogènes de celle-ci, et qui est choisi dans un groupe d'espèces appartenant au genre *Eimeria*, comprenant *E. acervulina*, *E. mivati*, *E. mitis*, *E. preacox*, *E. hagani*, *E. necatrix*, *E. maxima* et *E. burnetti*.

24. Composition d'immunogène d'*Eimeria*, comprenant une quantité immunologiquement efficace d'un ou plusieurs des immunogènes de groupe B de la revendication 23.

25. Utilisation de l'immunogène de la revendication 23 pour la préparation d'une composition utilisable pour l'immunisation de volailles contre la coccidiose provoquée par *Eimeria*.

Revendications pour l'Etat contractant suivant : ES

1. Procédé de préparation d'une molécule d'ADN recombinante, comprenant une séquence nucléotidique correspondant à l'ensemble ou à une partie de la séquence nucléotidique suivante:

```

T CTC GCC CCA ACT TTT TCC CCC GCG CTC CGC AGC AGC AGC AGC AGC AGC AGC AGC AGC
AGC AAA ATG GCA GAC CTC TTC AGC GGA CTC GTG GGC GGC GTC GTC GGC GCT GTT GGT
GCA GCA GAT TTG CCT GCG GAG GGC GAG AGG GCC CCC CGC CCC GCC CCC GGC ACT GGC
TGG ACT TGC TGC TGC AGC AAA CTG CAA GAA GGG GCC CGC GAG CTG GAG GGT TTT GTG
CAG CAG CTG AGT TTT GTT GCA GGG AAG CTG GCC TGC TGC CTG CGG GTG GGG GCG GAG
CAG CTG GCG CGC TGC GCT GCG GAG GGG CGG CTG CCC AGC AGC AGC AGC AGC AGC AGC
TGC TGC GCG CTG CTG CAG CTC GAG AAG CAG GAC CTC GAG CAG AGC CTC GAG GCC GGC
AAG CAG GGC GCG GAG TGC CTC TTG AGG AGC AGC AAA CTG GCC CTC GAG GCC CTC CTC
GAG GGG GCC CGC GTT GCA GCA ACG CGG GGT TTG CTG CTG GTC GAG AGC AGC AAA GAC
ACG GTG CTG CGC AGC ATT CCC CAC ACC CAG GAG AAG CTG GCC CAG GCC TAC AGT TCT
TTC CTG CGG GGC TAC CAG GGG GCA GCA GCG GGG AGG TCT CTG GGC TAC GGG GCC CTT
GCT GCT GCT TAC GGC CAG CAG CAG CAG CCC AGC AGC TAC GGG GCG CCC CCC GCC TCC
AGC CAG CAG CCC TCC GGC TTC TTC TGG TAG CCC TGC AGC AGC AGC AGC AGC AGC AGC
AGC AGC AGC AGC GCG GGC GGC AGC CGC GGC GGG GCC GGG GCG CCG CTG CAG CAA CAG
CAG CAG CCG nnn CGG CTA GCG CCG CGG AGC ACT CGC AGG GAA CTC CAC AGG CAG CCG
GAG AGC AGC AGG GAC GAG AAG CAG GTC ATG TAG CGC AGG CAG CAG CGC CAG CTG CAG
CAG CAG CAG CAG CAG CAG CAG CAG CAG CAG CAG CTC CTG CAC CG

```

qui code pour un immunogène de groupe B d'*Eimeria tenella*, comprenant l'obtention de ladite molécule d'ADN par la technique de l'ADN recombinant, et la combinaison de celle-ci avec un vecteur de clonage.

2. Procédé selon la revendication 1, dans lequel ladite séquence d'ADN est directement précédée par et fonctionnellement liée à une seconde séquence nucléotidique d'ADN, ce qui donne une molécule d'ADN fusionnée.

3. Procédé selon la revendication 2, dans lequel ladite seconde séquence d'ADN exprime une protéine CheY et le produit de nucléotides de liaison.

4. Procédé de préparation d'une molécule d'ADN recombinante comprenant une séquence nucléotidique capable d'être exprimée sous forme d'au moins une protéine manifestant l'immunogénicité d'un immunogène de groupe B d'*Eimeria tenella*, laquelle séquence est:

5 T CTC GCC CCA ACT TTT TCC CCC GCG CTC CGC AGC AGC AGC AGC AGC AGC AGC AGC AGC AGC
 AGC AAA ATG GCA GAC CTC TTC AGC GGA CTC GTG GGC GGC GTC GTC GGC GCT GTT GCT
 GCA GCA GAT TTG CCT GCG GAG GGC GAG AGG GCC CCC CGC CCC GCC CCC GGC ACT GCC
 10 TGG ACT TGC TGC TGC AGC AAA CTG CAA GAA GGG GCC CGC GAG CTG GAG GGT TTT GTG
 CAG CAG CTG AGT TTT GTT GCA GGG AAG CTG GCC TGC TGC CTG CGG GTG GGG GCG GAG
 CAG CTG GCG CGC TGC GCT GCG GAG GGG CGG CTG CCC AGC AGC AGC AGC AGC AGC AGC
 TGC TGC GCG CTG CTG CAG CTC GAG AAG CAG GAC CTC GAG CAG AGC CTC GAG GCC GGC
 15 AAG CAG GGC GCG GAG TGC CTC TTG AGG AGC AGC AAA CTG GCC CTC GAG GCC CTC CTC
 GAG GGG GCC CGC GTT GCA GCA ACG CGG GGT TTG CTG CTG GTC GAG AGC AGC AAA GAC
 ACG GTG CTG CGC AGC ATT CCC CAC ACC CAG GAG AAG CTG GCC CAG GCC TAC AGT TCT
 TTC CTG CGG GGC TAC CAG GGG GCA GCA GCG GGG AGG TCT CTG GGC TAC GGG GCC CCT
 20 GCT GCT GCT TAC GGC CAG CAG CAG CAG CCC AGC AGC TAC GGG GCG CCC CCC GCC TCC
 AGC CAG CAG CCC TCC GGC TTC TTC TGG TAG CCC TGC AGC AGC AGC AGC AGC AGC AGC
 AGC AGC AGC AGC GCG GGC GGC AGC CGC GGC GGG GCC GGG GCG CCG CTG CAG CAA CAG
 25 CAG CAG CCG nnn CGG CTA GCG CCG CGG AGC ACT CGC AGG GAA CTC CAC AGG CAG CGG
 GAG AGC AGC AGG GAC GAG AAG CAG GTC ATG TAG CGC AGG CAG CAG CGC CAG CTG CAG
 CAG CAG CAG CAG CAG CAG CAG CAG CAG CAG CAG CTC CTG CAC CG.

30 qui comprend l'obtention de ladite molécule d'ADN par la technique de l'ADN recombinant, et la combinaison de celle-ci avec un vecteur de clonage.

5. Procédé selon la revendication 4, dans lequel la séquence d'ADN est directement précédée par et fonctionnellement liée à une seconde séquence nucléotidique d'ADN, ce qui donne une molécule d'ADN fusionnée.

6. Procédé selon la revendication 5 dans lequel ladite seconde séquence d'ADN exprime une protéine CheY et le produit de nucléotides de liaison.

7. Procédé selon la revendication 1, 2 ou 5, dans lequel la séquence d'ADN est contenue dans un vecteur d'expression capable d'exprimer l'ensemble ou une partie de l'ADN immunogène, lorsqu'il est incorporé dans une cellule hôte.

8. Procédé de préparation d'un immunogène protéique d'*Eimeria tenella* recombinant, qui comprend l'expression de la molécule d'ADN de groupe B de la revendication 1, et toutes formes immunogènes de type sous-unité ou microhétérogènes de celle-ci.

9. Procédé de préparation d'un immunogène protéique de fusion d'*Eimeria tenella* recombinant, comprenant une protéine de fusion CheY liée par une protéine de liaison, qui est liée à une séquence d'acides aminés contenue dans un immunogène protéique d'*Eimeria tenella* de groupe B et des formes immunogènes de type sous-unité ou microhétérogènes de celle-ci, ledit procédé étant caractérisé par:

- l'obtention d'une molécule d'ADN codant pour la protéine;
- l'insertion de la molécule d'ADN dans un vecteur d'expression approprié;
- l'incorporation du vecteur d'expression dans une cellule hôte appropriée;
- la culture de la cellule hôte avec le vecteur d'expression, dans des conditions permettant l'expression de l'ADN et la production de la protéine; et
- la récupération de la protéine.

10. Procédé selon la revendication 9, dans lequel la séquence d'acides aminés contenue dans un immunogène protéique d'*Eimeria tenella* de groupe B comprend au moins la séquence d'acides aminés suivante:

5

10

Leu-Ala-Pro-Thr-Phe-Ser-Pro-Ala-Leu-Arg-Ser-Ser-Ser-Ser-Ser-Ser-Ser-

20 **30**

Ser-Ser-Lys-Met-Ala-Asp-Leu-Phe-Ser-Gly-Leu-Val-Gly-Gly-Val-Val-Gly-Ala-

40 **50**

Val-Ala-Ala-Ala-Asp-Leu-Pro-Ala-Glu-Gly-Glu-Arg-Ala-Pro-Arg-Pro-Ala-Pro-

15

60 70
 Gly-Thr-Ala-Trp-Thr-Cys-Cys-Cys-Ser-Lys-Leu-Gln-Glu-Gly-Ala-Arg-Glu-
 5
 80
 Leu-Glu-Gly-Phe-Val-Gln-Gln-Leu-Ser-Phe-Val-Ala-Gly-Lys-Leu-Ala-Cys-Cys-
 10
 90 100
 Leu-Arg-Val-Gly-Ala-Glu-Gln-Leu-Ala-Arg-Cys-Ala-Ala-Glu-Gly-Arg-Leu-Pro-
 15
 110 120
 Ser-Ser-Ser-Ser-Ser-Ser-Ser-Cys-Cys-Ala-Leu-Leu-Gln-Leu-Glu-Lys-Gln-Asp-
 20
 130 140
 Leu-Glu-Gln-Ser-Leu-Glu-Ala-Gly-Lys-Gln-Gly-Ala-Glu-Cys-Leu-Leu-Arg-Ser-
 25
 150 160
 Ser-Lys-Leu-Ala-Leu-Glu-Ala-Leu-Leu-Glu-Gly-Ala-Arg-Val-Ala-Ala-Thr-Agr-
 30
 170
 Gly-Leu-Leu-Leu-Val-Glu-Ser-Ser-Lys-Asp-Thr-Val-Leu-Arg-Ser-Ile-Pro-His-
 35
 180 190
 Thr-Gln-Glu-Lys-Leu-Ala-Gln-Ala-Tyr-Ser-Ser-Phe-Leu-Arg-Gly-Tyr-Gln-Gly-
 40
 200 210
 Ala-Ala-Ala-Gly-Arg-Ser-Leu-Gly-Tyr-Gly-Ala-Pro-Ala-Ala-Ala-Tyr-Gly-Gln-
 45
 220 230
 Gln-Gln-Gln-Pro-Ser-Ser-Tyr-Gly-Ala-Pro-Pro-Ala-Ser-Ser-Gln-Gln-Pro-Ser-
 Gly-Phe-Phe-Trp

et formes immunogènes de type sous-unité ou microhétérogènes de celle-ci.

11. Procédé selon la revendication 9, dans lequel la protéine de l'immunogène de groupe B d'*Eimeria tenella* est produite par la technique de l'ADN recombinant.
12. Procédé de production d'une protéine immunogène native purifiée d'*Eimeria tenella*, comportant une séquence d'acides aminés contenue dans la séquence d'acides aminés d'un immunogène protéique de groupe B d'*Eimeria tenella* et de toutes formes de type sous-unité ou microhétérogènes de celui-ci, qui comprend:
 - a. la désintégration d'ookystes sporulés d'*Eimeria tenella*, en présence d'inhibiteurs de protéase;

- b. la mise en contact des ookystes sporulés d'*Eimeria tenella*, désintégrés, avec un agent réducteur;
 c. la carboxyméthylation de la substance d'ookystes sporulés solubilisée, réduite; et
 d. la récupération séparément de l'immunogène protéique par chromatographie par exclusion de taille.

13. Procédé selon la revendication 12, dans lequel la séquence d'acides aminés comprend au moins la séquence d'acides aminés suivante:

10

Leu-Ala-Pro-Thr-Phe-Ser-Pro-Ala-Leu-Arg-Ser-Ser-Ser-Ser-Ser-Ser-Ser-

20

30

Ser-Ser-Lys-Met-Ala-Asp-Leu-Phe-Ser-Gly-Leu-Val-Gly-Gly-Val-Val-Gly-Ala-

40

50

Val-Ala-Ala-Ala-Asp-Leu-Pro-Ala-Glu-Gly-Glu-Arg-Ala-Pro-Arg-Pro-Ala-Pro-

60

70

Gly-Thr-Ala-Trp-Thr-Cys-Cys-Cys-Ser-Lys-Leu-Gln-Glu-Gly-Ala-Arg-Glu-

80

Leu-Glu-Gly-Phe-Val-Gln-Gln-Leu-Ser-Phe-Val-Ala-Gly-Lys-Leu-Ala-Cys-Cys-

90

100

Leu-Arg-Val-Gly-Ala-Glu-Gln-Leu-Ala-Arg-Cys-Ala-Ala-Glu-Gly-Arg-Leu-Pro-

110 120
 Ser-Ser-Ser-Ser-Ser-Ser-Ser-Cys-Cys-Ala-Leu-Leu-Gln-Leu-Glu-Lys-Gln-Asp-
 5 130 140
 Leu-Glu-Gln-Ser-Leu-Glu-Ala-Gly-Lys-Gln-Gly-Ala-Glu-Cys-Leu-Leu-Arg-Ser-
 10 150 160
 Ser-Lys-Leu-Ala-Leu-Glu-Ala-Leu-Leu-Glu-Gly-Ala-Arg-Val-Ala-Ala-Thr-Agr-
 15 170
 Gly-Leu-Leu-Leu-Val-Glu-Ser-Ser-Lys-Asp-Thr-Val-Leu-Arg-Ser-Ile-Pro-His-
 180 190
 Thr-Gln-Glu-Lys-Leu-Ala-Gln-Ala-Tyr-Ser-Ser-Phe-Leu-Arg-Gly-Tyr-Gln-Gly-
 20 200 210
 Ala-Ala-Ala-Gly-Arg-Ser-Leu-Gly-Tyr-Gly-Ala-Pro-Ala-Ala-Ala-Tyr-Gly-Gln-
 25 220 230
 Gln-Gln-Gln-Pro-Ser-Ser-Tyr-Gly-Ala-Pro-Pro-Ala-Ser-Ser-Gln-Gln-Pro-Ser-
 30
 Gly-Phe-Phe-Trp

et formes immunogènes de type sous-unité ou microhétérogènes de celle-ci.

- 35 14. Procédé de préparation d'une protéine immunogène native purifiée d'*Eimeria tenella*, comportant une séquence d'acides aminés contenue dans la séquence d'acides aminés mentionnée dans la revendication 13 et toutes formes de type sous-unité ou hétérogènes de celle-ci, qui comprend:
- 40 a. la désintégration d'ookystes d'*Eimeria tenella* sporulés, en présence d'inhibiteurs de protéase;
 b. la mise en contact des ookystes sporulés d'*Eimeria tenella*, désintégrés, avec un agent réducteur;
 c. la carboxyméthylation de la substance d'ookystes sporulés, solubilisée, réduite; et d. la récupération séparément de l'immunogène protéique par chromatographie d'immunoaffinité.
- 45 15. Procédé selon les revendications 9, 12 ou 14, dans lequel l'immunogène protéique de groupe B d'une espèce appartenant au genre *Eimeria* comprend un épitope ou déterminant antigénique correspondant à un immunogène d'*Eimeria tenella* de groupe B codé par l'ADN de la revendication 1, qui est identifié par un anticorps spécifique de l'immunogène de groupe B d'*Eimeria tenella*.
- 50 16. Procédé selon la revendication 15, dans lequel l'espèce est choisie dans un groupe d'espèces appartenant au genre *Eimeria*, comprenant *E. acervulina*, *E. mivati*, *E. mitis*, *E. preacox*, *E. hagani*, *E. necatrix*, *E. maxima* et *E. burnetti*.
- 55 17. Procédé selon la revendication 9, 12 ou 14, dans lequel l'immunogène comprend un épitope ou déterminant antigénique correspondant à la séquence d'acides aminés de la revendication 13 et à des formes immunogènes de type sous-unité ou microhétérogène de celui-ci, et qui est choisi dans un groupe d'espèces appartenant au genre *Eimeria*, comprenant *E. acervulina*, *E. mivati*, *E. mitis*, *E. preacox*, *E. hagani*, *E. necatrix*, *E. maxima* et *E. burnetti*.




FIG. 1




FIG. 2




FIG. 3





FIG. 4



S0216

FIG. 5




FIG. 6




FIG. 7




FIG. 8